

# **From cell surface to nucleus: Unraveling cancer metastasis and the role of nucleophosmin in breast cancer**

**Piia-Riitta Karhemo**

Research Programs Unit  
Translational Cancer Biology  
Institute of Biomedicine  
Faculty of Medicine  
&

Helsinki Graduate Program in Biotechnology and Molecular Biology  
University of Helsinki  
Finland

ACADEMIC DISSERTATION

*To be publicly discussed with the permission of the Faculty of Medicine,  
University of Helsinki, in lecture hall 3, Biomedicum Helsinki, Haartmaninkatu 8,  
Helsinki, on June 19<sup>th</sup> 2013 at 12 noon.*

Helsinki, 2013

**Supervised by**

PhD., Docent Pirjo Laakkonen  
K. Albin Johansson Senior Cancer Researcher, Finnish Cancer Organization  
Research Director  
Research Programs Unit, Translational Cancer Biology  
Biomedicum Helsinki  
University Helsinki  
Helsinki, Finland

**Reviewed by**

PhD., Docent Markku Varjosalo  
Head of Protein Research Group and Core Facility  
Institute of Biotechnology  
Biocenter 3  
University of Helsinki  
Helsinki, Finland

and

PhD., Professor Jorma Isola  
Institute of Biomedical Technology  
University of Tampere  
Tampere, Finland

**Opponent**

PhD., Professor Jonathan Sleeman  
Medical Faculty Mannheim  
University of Heidelberg  
Heidelberg, Germany

ISBN 978-952-10-8923-7 (paperback)  
ISBN 978-952-10-8924-4 (PDF)  
Helsinki University Printing House  
Helsinki 2013

# ABSTRACT

The most deadly aspect of cancer is its ability to spread from its original location to other sites in the body and grow as distant metastases. Formation of metastases is a multistep process and metastases can form even decades after the removal of the primary tumor. Cell surface proteins are known to play central roles in the adhesive contacts and molecular interactions between the tumor cell and the stroma during various stages of metastasis. In addition, they mediate important signals to intracellular proteins. As the detailed mechanisms of metastasis are still unclear, the aim of this thesis was to discover novel metastasis-associated cell surface proteins for further investigation.

This thesis established an optimized method for the isolation of biotinylated cell surface proteins for proteomic identification. This method was applied to compare the cell surface proteins isolated from an isogenic pair of human MDA-MB-435 cancer cell line with opposite metastatic phenotypes. We found 29 differentially expressed proteins and analyzed the molecular pathways they were involved in. Of these proteins expression of CD109 was shown to mark metastatic melanoma cells and invasive breast cancer cells.

Nucleophosmin (NPM) is a multitasking protein with both oncogenic and tumor suppressive functions. In our comparative proteomics analysis we discovered NPM to be expressed on the surface of the non-metastatic subclone of the MDA-MB-435 cells. We showed that NPM was detected at different localizations in the non-metastatic and metastatic cells most likely due to the expression of novel NPM splice variants discovered in this thesis work. In addition, we showed that expression level of NPM is one mechanism affecting its localization. In regards to patient prognosis, we revealed that high levels of NPM were expressed in the luminal epithelial cells of histologically normal breast tissue and that high levels independently associated with good prognosis in the luminal A breast cancer subtype. On the contrary, novel granular staining pattern and Threonine199 phosphorylation of NPM (NPMpThr199) correlated with aggressive characteristics, basal subtype and poor prognosis of human breast cancer. Moreover, NPMpThr199 associated with expression of a recently identified oncogene, cancerous inhibitor of protein phosphatase 2 (CIPA2).

In brief, this study provides several novel metastasis associated cell surface proteins for future investigation. By using breast cancer tumor microarrays from two large breast cancer patient cohorts and cellular models this thesis demonstrates for the first time, that different NPM forms play divergent and opposite roles in breast cancer.

# TIIVISTELMÄ

Syöpä on sairaus, jossa kehon solut jakaantuvat kontrolloimattomasti ja leviävät alkuperäisestä kasvupaikastaan elimistön muihin kudoksiin muodostaen niihin etäpesäkkeitä. Tämä johtaa potilaan kuolemaan. Syövän leviäminen ja etäpesäkkeiden muodostuminen on monivaiheinen prosessi, jonka kaikkia yksityiskohtia ei vielä tarkkaan tunneta. Koska syöpäsolut hyödyntävät solujen pinnalla ilmentyneitä proteiineja kommunikoidessaan ympäristönsä kanssa, näillä proteiineilla on tärkeä rooli syövän leviämisessä uusien kasvupaikkoihin. Väitöskirjatyöni tarkoituksena oli tunnistaa etäpesäkkeiden muodostumisen kannalta tärkeitä solun pintaproteiineja, joita ei ole aikaisemmin yhdistetty syövän leviämiseen.

Tutkimuksessa löysimme 29 proteiinia, joiden ilmentymisellä syöpäsolujen pinnalla oli yhteys solujen kykyyn muodostaa etäpesäkkeitä. Yksi löytämistämme proteiineista oli CD109, jota löytyi sekä etäpesäkkeistä eristetyistä melanoomasoluista että rintasyöpäsoluista, jotka pystyivät leviämään ympäröivään kudokseen. Jatkotutkimuksissa tulisi selvittää, voisiko CD109:n ilmentymistä käyttää syövän ennusteellisena tekijänä. Lisäksi tulisi selvittää, miten CD109:n läsnäolo syöpäsolujen pinnalla auttaa niitä leviämään elimistössä.

Tutkimuksen toisessa osassa löysimme merkittävän yhteyden nukleofosmiini-proteiinin (NPM) ilmentymistason ja potilaiden ennusteen välillä laajassa rintasyöpäaineistossa. Potilailla, joiden syöpäkasvaimissa NPM:n määrä oli vähentynyt, oli suurentunut riski etäpesäkkeiden muodostumiseen ja rintasyövästä aiheutuvaan kuolemaan. Lisäksi osoitimme solukokeissa NPM:n vähentävän rintasyöpäsolujen aggressiivisuutta. Nämä tuloksemme tukevat NPM:n tuumorisuppressiivista (kasvua estävää) toimintaa rintasyövässä. Toisaalta havaitsimme, että potilaiden huonoon ennusteeseen ja etäpesäkkeiden muodostumiseen vaikuttaa NPM:n treoniini-199-fosforylaatio. Havainto on merkittävä, sillä NPM:n toiminta on usein häiriintynyt kasvaimissa, mutta tämän häiriön taustalla vaikuttavia mekanismeja ei tarkkaan tunneta. Tutkimustuloksemme osoittavat sekä NPM:n ilmentymismäärän että sen fosforylaation vaikuttavan tämän proteiinin toimintaan rintasyövässä.

# CONTENTS

ABSTRACT .....	3
TIIVISTELMÄ .....	4
CONTENTS .....	5
LIST OF ORIGINAL PUBLICATIONS .....	7
ABBREVIATIONS.....	8
1 INTRODUCTION .....	10
2 REVIEW OF THE LITERATURE.....	11
2.1 Heterogeneity of cancer .....	11
2.1.1 Intrinsic breast cancer molecular subtypes .....	12
2.2 Metastatic dissemination of cancer.....	14
2.2.1 The metastatic cascade.....	15
2.2.2 Organ specific colonization and microenvironment.....	18
2.2.3 Dormancy of tumor cells .....	19
2.2.4 When tumor cells acquire their metastatic capability? .....	21
2.3 Subclones of MDA-MB-435 cancer cell line as a metastasis model .....	23
2.3.1 Origin of the parental MDA-MB-435 cell line? .....	23
2.3.2 Cloning and characterization of the non-metastatic and metastatic subclones of the MDA-MB-435.....	24
2.3.3 Identification of metastasis related cell surface proteins .....	24
2.4 Nucleophosmin; oncogene, tumor suppressor or both? .....	26
2.4.1 Discovery of NPM as nucleolar protein involved in ribosome assembly and transport .....	31
2.4.2 NPM as a chaperone and its role in transcriptional regulation.....	31
2.4.3 Molecular pathways regulated by NPM: RB, ARF-p53- Mdm2 pathway and c-Myc.....	33
2.4.4 Role of NPM in centrosome duplication.....	36
2.4.5 NPM in human tumors and in mice .....	37
3 AIMS OF THE STUDY .....	39
4 MATERIALS AND METHODS .....	40
4.1 Cell culture (2D and 3D), cell treatments, cell proliferation and soft agar assays .....	42

4.2	Cell surface protein isolation.....	43
4.3	Mass spectrometry, protein identification, network and pathway analyses.....	44
4.4	Immunoprecipitation and immunoblotting.....	45
4.5	Immunohistochemistry and immunofluorescence analyses .....	45
4.6	Patients and classification of breast cancer subtypes in the TMAs .....	46
4.7	Statistical analyses .....	46
5	RESULTS AND DISCUSSION .....	47
5.1	Comparative analysis of metastasis and dormancy associated cell surface proteins (I and unpublished data) .....	47
5.1.1	Optimized method for isolation of biotinylated cell surface proteins (I) .....	48
5.1.2	Cell surface protein changes in the metastatic cells (I, unpublished data) .....	49
5.1.3	Validation of the differential cell surface expression of selected proteins (I, unpublished data) .....	52
5.1.4	The overexpression of cd109 and ITGA6 marks cells derived from melanoma metastasis (I) .....	54
5.1.5	Interactome analysis indicates a role for a tetraspanin protein complex in metastasis (I) .....	56
5.2	Nucleophosmin (NPM/B23) has multiple divergent roles in breast cancer (II, III) .....	58
5.2.1	Discovery and characterization of novel NPM splice variants B23.3 and B23.4 (III) .....	58
5.2.2	Analysis of the prognostic significance of different NPM forms in breast cancer (II, III) .....	60
5.2.3	NPM expression levels affected its localization (II) .....	62
5.2.4	Tumor suppressive role for NPM in breast cancer (II) .....	63
5.2.5	Granular staining pattern as indicator of metastatic disease (II).....	64
5.2.6	Thr199 phosphorylation of NPM as an indicator of recurrence (III) .....	64
5.2.7	Thr199 phosphorylation of NPM associated with aggressive breast cancers and CIP2A expression (III).....	65
5.2.8	NPM modifications and metastasis? .....	66
6	CONCLUSIONS AND FUTURE PROSPECTS .....	68
7	ACKNOWLEDGEMENTS .....	69
8	REFERENCES.....	71

# LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following original publications, which are referred to in the text by their Roman numerals. Unpublished material is also presented.

- I. Piia-Riitta Karhemo, Suvi Ravela, Marko Laakso, Ilja Ritamo, Olga Tatti, Selina Mäkinen, Steve Goodison, Ulf-Håkan Stenman, Erkki Hölttä, Sampsa Hautaniemi, Leena Valmu, Kaisa Lehti, Pirjo Laakkonen. An Optimized Isolation Of Biotinylated Cell Surface Proteins Reveals Novel Players In Cancer Metastasis, J Proteomics. 2012 Dec 21;77:87-100.  
doi: 10.1016/j.jprot.2012.07.009
- II. Piia-Riitta Karhemo, Antti Rivinoja, Johan Lundin, Maija Hyvönen, Anastasiya Chernenko, Johanna Lammi, Harri Sihto, Mikael Lundin, Päivi Heikkilä, Heikki Joensuu, Petri Bono, Pirjo Laakkonen. An Extensive Tumor Array Analysis Supports Tumor Suppressive Role for Nucleophosmin in Breast Cancer, Am J Pathol. 2011 August; 179(2): 1004–1014. doi: 10.1016/j.ajpath.2011.04.009
- III. Piia-Riitta Karhemo, Harri Sihto, Anni Laine, Antti Rivinoja, Petri Bono, Henna Moore, Päivi Rajahalme, Marikki Laiho, Jukka Westermarck, Heikki Joensuu and Pirjo Laakkonen. Novel players in oncogenicity of NPM? (manuscript)

The original publications are reproduced with the permission of the publishers.

# ABBREVIATIONS

AML	acute myeloid leukemia
ARF	p14ARF
ATP	Adenosine-5'-triphosphate
BLBC	basal like breast cancer
CIP2A	cancerous inhibitor of protein phosphatase 2
CD109	cluster of differentiation 109
CDK	cyclin-dependent kinase
CK2	casein kinase 2
CK5/6	cytokeratin-5/6
CSCs	cancer stem cells
CTCs	circulating tumor cells
DTCs	disseminated tumor cells
ECM	extracellular matrix
ECGFP	enhanced cyan green fluorescent protein
EGFR	epidermal growth factor receptor
EMT	epithelial to mesenchymal transition
ER	estrogen receptor
GRK5	G protein-coupled receptor kinase 5
HER1	human epidermal growth factor receptor-1
HER2	human epidermal growth factor receptor-2
IGSF8	immunoglobulin superfamily member 8
ITGA6	integrin- $\alpha$ 6
ITGB1	Integrin- $\beta$ 1
KEGG	Kyoto Encyclopedia of Genes and Genomes
LC-MS/MS	liquid chromatography coupled to tandem mass spectrometry
MDM2	murine double minute 2
MEFs	mouse embryonic fibroblasts
NPM	nucleophosmin
pThr199	threonine 199 phosphorylation
PP1	protein phosphatase 1



PR	progesterone receptor
PROCR	endothelial protein C receptor
PTGFRN	prostaglandin receptor negative regulator
PTP	protein tyrosine phosphatase
PTPRF	receptor-type tyrosine-protein phosphatase F
RB	retinoblastoma protein
rRNA	ribosomal RNA
SENP	SUMO1/sentrin/SMT3 specific peptidase
SSR4	Translocon associated protein delta subunit
SUMO	Small ubiquitin-like modifier
TICs	tumor initiating cells
TMA	tissue microarray
TME	tumor microenvironment
TNBC	triple negative breast cancer
VAMPA	membrane protein-associated protein A
2D	two dimensional
3D	three dimensional

# 1 INTRODUCTION

Cancer comprises a large group of diseases caused by deregulated cell growth and resistance to cell death in any part of the body. Every fourth person in Finland is diagnosed with cancer at some point of their life (Cancer Statistics at Finnish Cancer Registry) and according to the World Health Organizations GLOBOCAN 2008 project 7.6 million annual cancer deaths are reported worldwide (Ferlay et al., 2011). The most deadly aspect of cancer is its ability to spread from its original location to other sites in the body in a process referred to as metastasis. Metastasis is a multistep process, details of which are not yet fully understood. Therefore, there is a need to identify novel proteins and molecular pathways that are involved in the regulation of cancer metastasis to better understand and disrupt the process.

Proteins perform many important cellular functions. Mutations in protein encoding genes can modify protein function in many diseases such as cancer (Futreal, 2004). To understand how cancer cell acquires a metastatic phenotype, it is important to know the quantities of different proteins in normal and malignant cells and understand how these quantities change. Proteomics techniques are used to study proteins present in a cell, tissue or an organism at given time and can be applied to find changes in proteins and their quantities between different conditions or cell types like metastatic and non-metastatic tumor cells.

Tissue microarrays (TMAs) allow analysis of tissue specimens at nucleic acid or protein level (Kononen et al., 1998; Avninder et al., 2008). These have vastly facilitated the clinical validation of molecular discoveries made with the aid of genomics and proteomics methods. TMAs are constructed from archival formalin-fixed paraffin embedded tissue and can contain patient follow-up data to help classify the clinical significance of the findings. In this thesis work, proteomics was used to find novel proteins that might play important roles in cancer metastasis. Furthermore, the role of one of the identified proteins, nucleophosmin (NPM), was analyzed in breast cancer by using TMAs as well as cell biological and biochemical methods.

## 2 REVIEW OF THE LITERATURE

### 2.1 Heterogeneity of cancer

Cancer refers to a collection of heterogeneous malignancies in various locations in the body. Cancer is caused by uncontrolled cell growth and resistance to cell death. Slow accumulation of alterations in proto-oncogenes, tumor-suppressor genes, DNA-repair genes and microRNA genes together with epigenetic changes in one cell or a small group of cells is considered to lead to cancer development with varying times depending on the tumor type. Recent, debatable theories suggest that not all cancer cells in a tumor are alike and that only so called cancer stem cells (CSCs) or tumor-initiating cells (TICs) would be able to maintain the tumor by possessing the ability to self-renew and proliferate. Other tumor cells would differentiate into cells that constitute the bulk of the tumor mass (Reya, 2001; Zhou, 2009). The neoplastic cancer cells harboring genetic alterations do not manifest the disease alone but form organ-like structures together with the tumor microenvironment (TME), which is composed of different types of normal stromal cells and the extracellular matrix (ECM). Consequently, cancer formation depends on both cancer cell-intrinsic pathways and cancer cell-extrinsic pathways (Hanahan, 2012).

Cancers are categorized into different types depending on their tissue of origin. Carcinomas like lung, breast and colon cancer originate from epithelial tissues and represent the most common cancer type. Non-epithelial cancers can be divided into i) sarcomas which originate from mesenchymal cells, ii) hematological cancers (leukemias and lymphomas), which originate from hematopoietic cells and iii) neuronal cancers (gliomas, glioblastomas, neuroblastomas, schwannomas, medulloblastomas) which originate from various components of the central and peripheral nervous system. Recent gene array technologies have, however, revealed a heterogeneity in tumors appearing in the same organ i.e. lung (West et al., 2012), skin (Vidwans et al., 2011) or breast (Perou et al., 2000; Sorlie et al., 2001; Sorlie et al., 2003; Hu et al., 2006) . This information can be used to separate the breast, skin or

lung tumors into several distinct molecular subtypes and in the future help to develop individual treatment guidelines for the different subtypes.

### **2.1.1 Intrinsic breast cancer molecular subtypes**

Breast cancer is a heterogeneous group of diseases in terms of histology, therapeutic response, metastatic dissemination, and patient outcomes and it has recently been divided into the following intrinsic biological subtypes (Perou et al., 2000; Sorlie et al., 2001; Sorlie et al., 2003; Hu et al., 2006): Luminal A, Luminal B, basal-like breast cancer (BLBC), human epidermal growth factor receptor-2-enriched (HER2-enriched) and normal-like. Of these, Luminal A and B are positive for estrogen receptor (ER) while BLBC and HER2-enriched tumors are ER negative (Goldhirsch et al., 2011). Luminal A and B subtypes differ from each other in their HER2 expression and/or proliferation index so that luminal A tumors are HER2-negative and luminal B tumors HER2-positive. Recently, a new intrinsic, claudin low subtype of breast cancer, was also suggested (Prat et al., 2010).

Gene expression profiling is not used in clinical practice to classify tumors. Therefore, the gene array-based intrinsic subtypes have been evaluated in immunohistochemistry by using antibodies against common markers determining the subtypes: ER, progesterone receptor (PR) and HER2. In addition, epidermal growth factor receptor (EGFR) (Carey et al., 2006), cytokeratin-5/6 (CK5/6) (Carey et al., 2006; Nielsen et al., 2004; Blows et al., 2010), and markers like human epidermal growth factor receptor-1 (HER1) (Nielsen et al., 2004) or Ki67 (Hugh et al., 2009; Cheang et al., 2009) have been used to classify the basal subtype, depending on the study.

The molecular subtypes differ in their mutation status for the tumor suppressor protein p53. Only about 12-15% of luminal A tumors harbor p53 mutations while function of p53 is lost by mutation or other means in most of the BLBCs (Carey et al., 2006; Cancer Genome Atlas Network, 2012; Dumay et al., 2013). In addition to molecular subtypes, breast cancers can be classified as triple negative (TNBC), which shows negative staining for HER2,

ER and PR (Reis-Filho and Tutt, 2008). TNBCs comprise of various kinds of tumors, but majority of them are BLBCs (Carey et al., 2010). The tumor suppressor RB is commonly affected in TNBC and BLBC (Gauthier et al., 2007; Herschkowitz et al., 2008; Subhawong et al., 2009). In addition, most *BRCA1* mutant breast cancers are both triple negative and basal-like (Turner and Reis-Filho, 2006; Atchley et al., 2008; Hartman et al., 2012).

The prognosis of breast cancer patients is generally favorable and mortality has declined due to early detection and improved adjuvant therapies (Schopper and de Wolf, 2009). However, the metastatic dissemination of breast cancer to other organs is not uncommon and women with advanced disease still have a median survival time of only approximately two years (Largillier et al., 2008; Anderson et al., 2000). Currently, most of the breast cancer patients are treated with adjuvant therapy because of the lack of proper prognostic and predictive markers of metastasis. Novel markers of metastasis are needed to help clinicians to select the estimated 40% of patients that will benefit from the adjuvant therapy. In addition, the quality of life would increase for the patients that can be cured by local treatment only since they would not have to needlessly suffer from the side effects of the adjuvant therapy (Weigelt et al., 2005).

The molecular differences in breast cancer subtypes result in distinct clinical outcomes and responses to treatment; in general, the luminal A tumors associate with favorable and the BLBC and HER2-enriched tumors with poor prognosis (Carey et al., 2006; Voduc et al., 2010; Dawood et al., 2011; Arvold et al., 2011; Sorlie et al., 2001). The subtypes also have distinct preference for their metastatic sites. Luminal A cancers metastasize first to bone, HER2-enriched cancers to liver and lung and basal cancers to liver and brain (Sihto et al., 2011; Smid et al., 2008).

The biological mechanisms for breast cancer heterogeneity are mainly unknown. Possible explanations include distinct cell of origin, like CSCs or progenitor cells and tumor subtype-specific genetic events. These two mechanisms are not necessarily mutually exclusive. Two major epithelial cell populations are found in the mammary gland; the inner luminal epithelial cells and the outer (basal) myoepithelial cells, embedded in a stromal matrix.

These two populations can be divided in further sub-populations. For example, the luminal layer is a mixed population of ER positive and ER negative cells. Functionally, the mammary epithelial cells can be classified as stem, progenitor and differentiated cells. Luminal compartment contains majority of the progenitor cells while the stem cell activity is mainly found in the basal layer (Molyneux and Smalley, 2011). Mouse models (Ginestier et al., 2012) and the presence of these different cell populations in the mammary gland supports the hypothesis that breast cancer heterogeneity would result from different molecular changes occurring in different cell types (Dontu et al., 2003). Recent research indicates that BRCA1-associated breast cancers and potentially also non-familial BLBC and TNBC would originate from the luminal ER negative progenitors (Molyneux and Smalley, 2011).

## **2.2 Metastatic dissemination of cancer**


One of the hallmarks in cancer progression is the acquisition of an invasive phenotype that allows cancer cells to spread to distant sites in the body and form metastatic lesions that are resistant to many cancer treatments (Hanahan and Weinberg, 2000; Hanahan and Weinberg, 2011). The term “metastasis” was invented in 1829 by a French gynecologist Recamier (Recamier, 1829) to describe the spread of cancer from its original growth sites to other parts in the body. Even today, the metastatic dissemination, rather than the primary tumor, is responsible for 90% of cancer deaths making metastasis the most serious challenge for cancer treatment (Gupta and Massague, 2006). The formation of metastasis depends on multiple steps the details of which are still poorly understood. In addition to the tumor cells TME or stroma participates in tumor progression, metastasis and response to treatment (Sleeman et al., 2012; Joyce and Pollard, 2009). In breast cancer, ECM and numerous stromal cell types, including endothelial and immune cells, fibroblasts, and adipocytes make up the primary tumor TME (Place et al., 2011).

Proteins and multiprotein complexes at the cell surface play important roles in sensing the environment, the signaling and the adhesive contacts between tumor and stromal cells during different stages of the metastatic dissemination (Karhemo et al., 2012; Place et al., 2011; Bendas and Borsig, 2012). In addition, tumor cells use their cell surface proteins to interact with platelets, leukocytes, and soluble components during the establishment of metastatic lesions (Bendas and Borsig, 2012). In the following paragraphs some examples of known cell surface proteins affecting the different parts of the metastasis process are given.

### **2.2.1 The metastatic cascade**

Cancer cells must complete a set of well-defined, interrelated steps, globally referred to as the metastatic cascade, to develop clinically detectable metastases. Cancer cells must detach from the primary tumor, invade and survive in the lymphatics and/or blood vessels to be transported to distant organs where they must adhere, extravasate, and proliferate to form a metastatic lesion (Valastyan and Weinberg, 2011; Fidler, 2003) (Figure 1).

Due to the complexity of the metastasis cascade, metastasis is an inefficient process and can fail at any step (Chambers et al., 2002; Mehlen and Puisieux, 2006; Fidler, 1970). The entry of tumor cells into the circulation is common and more than a million cells per gram of tumor can be shed daily (Butler and Gullino, 1975). However, only a fraction of the shed tumor cells actually survive in the circulation as circulating tumor cells (CTCs) or in bone marrow as disseminated tumor cells (DTCs) (Ross and Slodkowska, 2009; Mehlen and Puisieux, 2006).



**Figure 1.** The Invasion-Metastasis Cascade. To be able to grow as a distant metastasis, cancer cell need to go through a series of different steps termed the invasion-metastasis cascade. The cell needs to invade the surrounding tissue (local invasion), enter the circulation (intravasation), survive in the circulation to be transported to distant sites. A cancer cell needs to be able to attach to the capillary wall at the distant organ, extravasate to form micrometastasis. Finally, to grow as a full-blown metastasis, cancer cell needs to avoid metastatic dormancy and proliferate at the distant site. (Adapted and modified from (Valastyan and Weinberg, 2011)).



Recently, it was suggested that this complex metastatic cascade could be simplified into two major phases: i) physical translocation from the primary tumor to the distant organ (contains the invasion, survival in the circulation and transport to the distant site) and ii) colonization at the distant site (Chaffer and Weinberg, 2011). The molecular mechanisms underlying the first phase are quite well resolved. During embryogenesis cells lose their epithelial characteristics and gain mesenchymal properties in a process entitled epithelial to mesenchymal transition (EMT). EMT is considered to play a role at least in the first phase of metastasis by changing the adhesive properties of tumor cells and promoting their motility, thereby increasing their invasiveness (Berx et al., 2007). However, the functional contribution of EMT to metastasis in patients is still debated (Sleeman et al., 2012).

At the cell surface perspective, deregulated expression of many cell surface ECM remodeling enzymes like heparanases and matrix metalloproteinases is a common event in human cancers to help cancer cells in their invasion (Lu et al., 2011). In most tumors cell surface proteins and protein complexes like integrins, syndecans, dystroglycans, immunoglobulin superfamily cell adhesion proteins, cadherins, and hyaluronan binding proteins (hyaladherins) like CD44 participate in cell–tissue interaction and migration during invasion (Gritsenko et al., 2012). Recently, microvesicles, small membrane-enclosed sacs, shed from tumor cells have been shown to participate in the regulation of the ECM invasion and evasion of the immune response (Muralidharan-Chari et al., 2010).

The mechanisms playing a role in the second phase of metastasis, colonization still remains largely unknown. In experimental models survival and proliferation at the secondary site, have been shown to be highly inefficient (Allan et al., 2006). Clinically, it would be of utmost importance to better understand the second phase to be able to treat patients who have already developed metastasis at the time of diagnosis (Chaffer and Weinberg, 2011). Most likely, cell surface proteins play a major role also in this phase.

The models mentioned above describe metastasis as a unidirectional process, where cancer cells from the primary tumor seed metastasis in distant sites. Interestingly, based on recent experimental models a new

metastasis concept, termed self-seeding, has been proposed. This paradigm considers metastasis as a multidirectional process whereby cancer cells can seed distant sites as well as the primary tumor itself (Norton and Massague, 2006; Comen et al., 2011).

### **2.2.2 Organ specific colonization and microenvironment**

The distribution of full-blown metastases to different organs is not random and different tumor types disseminate and form metastatic lesions in a different set of organs (Auerbach et al., 1987; Johnson et al., 1991; Nguyen et al., 2009). Already in 1889 Stephen Paget proposed, based on his analysis of autopsy records from 735 breast cancer patients, that DTCs, or “seeds,” would only colonize organ microenvironments, or “soils,” that would be compatible with the growth of the DTCs (Paget, 1889). This so called seed and soil hypothesis suggested that outcome of metastasis depends on the interactions between the tumor cells and host tissue, a fact that is currently emerging as a critical determinant of metastasis (Lorusso and Ruegg, 2012).

The seed and soil hypothesis was challenged by Ewing by stating that organ specificity is accounted by mechanical forces and circulatory patterns between the primary tumor and the secondary site (Ewing, 1928). Later, Fidler and coworkers (Fidler and Kripke, 1977; Hart and Fidler, 1980) revealed that although CTCs in the vasculature traffic through all organs, metastases selectively develop in organs with suitable environment. In support of this, breast cancer frequently metastasizes to the lungs, bones and liver (Largillier et al., 2008) and brain (Palmieri et al., 2006), which do not have a direct circulatory connection to breast tissue (Lorusso and Ruegg, 2012).

Mechanistically, networking of cytokines and chemokines expressed in the target tissue and interacting with their cognate receptors expressed on the surface of the tumor cells is involved in organ specificity highlighting the important role of cell surface proteins in this process. As examples, production of osteoclast-activating factors such as pTHRp, Il-11, Il-6, tumor necrosis factor- $\alpha$  (TNF $\alpha$ ), and granulocyte–macrophage colony stimulating

factor (GM-CSF) is required for the ability of breast cancer cells to form bone metastases (Nguyen et al., 2009).

Recent evidence from cell line and animal studies suggests that specific cell surface adhesion molecules on tumor cells and their receptors on the lung endothelium mediate breast cancer cell adhesion and extravasation in the lung (Lu and Kang, 2007). In breast cancer, the CXCR4 and CCR7 receptors, expressed on breast cancer cell surface and their ligands CXCL12 and CCL21 on the organs, might play a role in cancer cell arrest and migration into secondary organs (Muller et al., 2001; Wang et al., 2006). Metadherin, which is overexpressed on the surface of metastatic breast cancer cells, mediates targeting of tumor cells specifically to the lung, but not to other organs through binding to an unknown receptor expressed in lung endothelium (Brown and Ruoslahti, 2004). In addition, interaction of cell surface fibronectin with a dipeptidyl peptidase IV (DPP IV, (Cheng et al., 1998) and cell surface expression of  $\alpha 6 \beta 4$  integrin and its adhesion to human CLCA2 protein (Abdel-Ghany et al., 2001) were shown to participate in the lung metastasis of breast cancer.

As described, the successful engraftment and growth of cancer cells in distant organs depends on a receptive microenvironment. Recent evidence points out that growth factors and other molecules secreted by the primary tumor could prime certain tissues for tumor cell engraftment by forming a pre-metastatic niche (Psaila and Lyden, 2009; Psaila et al., 2006). The location of these pre-metastatic niches would subsequently determine the organs in which metastases will form (Sleeman and Cremers, 2007).

### **2.2.3 Dormancy of tumor cells**

Metastases can occur after long latency periods that range from years to decades after the primary treatment (Chaffer and Weinberg, 2011). For example, about 45% of breast cancer patients will relapse and develop distant metastases years or decades after the diagnosis (Karrison et al., 1999). This prolonged time observed for the development of distant, metastatic disease

indicates a period of dormancy before DTCs are able to grow into clinically relevant metastases (Castano et al., 2011).

In an experimental model, dormant tumor cells have been observed in metastasis free organs of animals carrying spontaneously metastatic primary tumors (Suzuki et al., 2006) suggesting that dormancy might also affect organ specific colonization so that in the hostile environments DTCs undergo dormancy while in a receptive environment the same cells are able to grow into macrometastases.

Concept of tumor cell dormancy was introduced already in 1934 by an Australian pathologist Rupert Willis (Willis, 1934). Despite the early observation of cancer dormancy, not much is known about cancer cells during this period of dormancy and what awakens them even though this is clinically a very important question (Uhr and Pantel, 2011). Dormant cells may be reactivated by modification of their microenvironment (Barkan et al., 2010; Barkan et al., 2010) or by loss of metastasis suppressor genes, which are defined by their ability to inhibit overt metastasis in a secondary organ without affecting tumor growth at the primary site (Horak et al., 2008).

Two forms of dormancy, which are not mutually exclusive, have been suggested (Castano et al., 2011). Dormancy might be accounted for by a mitotic arrest of tumor cells (Naumov et al., 2002; Goodison et al., 2003; Muller et al., 2005), which is called cellular dormancy (Castano et al., 2011). In another form of dormancy, called tumor dormancy, the rate of cell death counterbalances the rate of cell proliferation within a tumor mass (Meng et al., 2004; Naumov et al., 2006). However, the mechanisms controlling the size of the tumor cell population are unknown (Meng et al., 2004) but it might be kept constant by some of the same mechanisms that control the size of normal organs (Uhr and Pantel, 2011).

The molecular mechanisms of dormancy have mostly been studied in experimental models. Dormant state might be regulated by a crosstalk between tumor cell surface proteins and the ECM at the secondary sites (Barkan et al., 2010). As examples, cell surface urokinase receptor (Aguirre Ghiso et al., 1999; Liu et al., 2002; Heiss et al., 1995; Aguirre-Ghiso et al., 2001) and integrin  $\beta$ 1-focal adhesion kinase (FAK) signaling axis has been

shown to regulate metastatic dormancy in three dimensional *in vitro* model (Shibue and Weinberg, 2009; Pontier and Muller, 2008). Maintenance of the dormant state might also be affected by failure of angiogenesis (Goss and Chambers, 2010). Recently, Kim et al. revealed a dormancy signature in breast cancer and showed that ER-positive tumors, which generally associate with favorable prognosis and carry a dormancy signature, are likely to undergo prolonged dormancy before resuming metastatic growth (Kim et al., 2012).

#### **2.2.4 When tumor cells acquire their metastatic capability?**

To understand the properties of metastatic tumor cells and target the process, it is important to resolve, when the tumor cells acquire their full metastatic capability and are able to go through both phases of metastasis. Two basic models have been suggested to represent the timing of the metastatic cascade during tumor progression. The prevailing, so called linear progression model, suggests that a rare subpopulation of tumor cells in the primary tumor acquires a full malignant phenotype via genomic alterations at late stages of primary tumor development. These fully malignant cells would have all the properties required for the formation of metastatic lesions (Fidler and Kripke, 1977; Klein, 2009; Poste and Fidler, 1980). The association of large tumor size with higher frequency of metastases (Koscielny et al., 1984), the correlation between primary tumor size and risk of lymph node and distant metastasis (Comen et al., 2011), the curative effect of surgery on smaller lesions (Klein, 2009) and the variable metastatic capability of different murine B16 melanoma cell clones (Fidler and Kripke, 1977) support the linear progression model.

A recent, parallel model of metastasis questions the linear progression model and proposes that metastases arise from DTCs, which do not necessarily disseminate near the end of primary tumor development and acquire their fully metastatic phenotype independent of the primary tumor (Klein, 2009). Quantitation of human cancer growth rates demonstrates that the metastatic lesions must have been initiated long before the diagnosis of

the primary tumor (Collins et al., 1956; Friberg and Mattson, 1997; Klein, 2009). Furthermore, the long interval between diagnosis and relapse with metastatic, distant disease in breast cancer, e.g. dormancy, supports the parallel progression model (Nguyen et al., 2009). In addition, genetic comparison of breast cancer metastases to their matched primary tumors supports the parallel progression and early onset of metastases (Sleeman et al., 2012; Kuukasjarvi et al., 1997; Torres et al., 2007; Santos et al., 2008). In addition, recent studies show that breast cancer derived DTCs display fewer genetic alterations than their corresponding primary tumors (Schardt et al., 2005; Schmidt-Kittler et al., 2003) and these alterations do not resemble those detected in the corresponding primary tumors (Lorusso and Ruegg, 2012; Schmidt-Kittler et al., 2003; Braun et al., 2005). However, DTCs are currently isolated for analysis by the aid of certain epithelial markers, which might not be expressed by all DTCs such as tumor cells undergoing EMT hampering the reliable comparison of all DTCs to primary tumors (Sleeman et al., 2012).

It has also been suggested that particular early oncogenic events that drive primary tumor growth might give cancer cells their propensity to metastasize as opposed to the theory that metastasis arises from rare cells accumulating metastasis specific genomic alterations in time (Bernards and Weinberg, 2002; Ramaswamy et al., 2003). Another explanation for the metastatic paradox was revealed from metastatic studies in different genetic background in mice. These preliminary studies indicate that the host genetic background has a significant role in determining the metastatic potential early in oncogenesis (Hunter et al., 2003).

## **2.3 Subclones of MDA-MB-435 cancer cell line as a metastasis model**

### **2.3.1 Origin of the parental MDA-MB-435 cell line?**

The MDA-MB-435 cell line was created from malignant cells in a pleural effusion of a 31-year old Caucasian woman with breast cancer. Patient had an extensive infiltrating breast carcinoma and two of the eight axillary lymph nodes contained breast cancer cells. She died one year after her diagnosis because of a metastatic disease (Cailleau et al., 1978; Brinkley et al., 1980).

The origin of the cell line has later been questioned. The microarray data and karyotyping show that the cell line has a gene expression pattern most compatible with melanocyte origin and identical to the M14 melanoma cells that were used as a feeder cell line during the establishment of the MDA-MB-435 cell line (Ross et al., 2000; Rae et al., 2004; Rae et al., 2007). In support of the melanoma origin, the MDA-MB-435 cells were shown to express RXRG, TYR, ACP5, and DCP genes, which are commonly transcribed in melanocytes but not in various commonly used breast cancer cell lines (Ellison et al., 2002).

However, MDA-MB-435 cells can be induced to express breast differentiation markers and secrete milk lipids (Sellappan et al., 2004). They also express a number of breast and epithelial cell specific proteins together with melanocytic features, most likely due to lineage infidelity (Sellappan et al., 2004; Nerlich and Bachmeier, 2013). It is possible that the MDA-MB-435 cells represent undifferentiated breast cancer and express melanocytic differentiation markers since primary breast tumors have been shown to express melanocyte related genes (Montel et al., 2009). Furthermore, based on karyotype and allelotype, the MDA-MB-435 cells are of female origin and cannot therefore be classified as M14 melanoma, which originate from a male patient (Chambers, 2009; Hollestelle and Schutte, 2009).

In respect to the breast cancer molecular subtypes, MDA-MB-435 cell line has been classified to represent the basal subtype (Neve et al., 2006; Chavez

et al., 2010). It contains a wild-type BRAC1 (Elstrodt et al., 2006) and a mutant p53 (Hollestelle et al., 2010; O'Connor et al., 1997).

### **2.3.2 Cloning and characterization of the non-metastatic and metastatic subclones of the MDA-MB-435**

Limiting dilution technique, with direct microscopic monitoring of monocellular origin, has been used to create a pair of isogenic clones of the MDA-MB-435 cell line. Screening for the metastatic ability in athymic mice revealed that these clones significantly differ in their metastatic capability (Urquidi et al., 2002). Both the metastatic and non-metastatic cells are able to reach the lungs of tumor-bearing mice thus capable of going through the first stage of metastasis, the physical translocation, while only the metastatic cells can perform the second phase of metastasis, colonization, in lungs and form full-blown metastatic lesions (Goodison et al., 2003). In addition, metastatic cells that formed lung metastases could be observed in a dormant state in other organs (Suzuki et al., 2006). Thus, these cell lines enable the comparative investigation of cellular and molecular events necessary for the second phase of metastasis and for the maintenance and subsequent release from dormancy at the secondary sites in a stable and isogenic model.

### **2.3.3 Identification of metastasis related cell surface proteins**

As described earlier, the details and molecular mechanisms of metastasis are not fully resolved. At the late stages of metastasis blood flow and other mechanical factors influence the delivery of cancer cells to specific organs, whereas molecular interactions between the cancer cells and the organ influence the probability that the cells will proliferate and grow as a metastatic lesion at the new site. Cell surface proteins, the proteins protruding from the plasma membrane into the extracellular space, are important mediators of these interactions (Place et al., 2011; Bendas and Borsig, 2012; Karhemo et al., 2012). Cell surface molecules also represent two-thirds of the current protein-based drug targets (Hopkins and Groom,



2002; Overington et al., 2006). Some, but not all, cell surface proteins can be classified as plasma membrane proteins. For example, ligands bound to their surface receptors can be regarded only as cell surface proteins because they lack direct contact with the plasma membrane.

Heterogeneity of tumors and presence of stromal cells within tumors hamper the search for cancer cell specific metastasis-associated proteins (Hondermarck et al., 2008). Large-scale analysis of cell surface proteins is hindered by the poor solubility of hydrophobic, integral membrane proteins. Cell surface and membrane proteins are also of low abundant and difficult to detect without enrichment or fractionation. Most cells can be removed from tissues, but this is difficult to perform without perturbing the cell surface (Leth-Larsen et al., 2010). For these reasons, it is difficult to study these proteins *in vivo* at tissue level.

The use of isogenic cell lines differing in their metastatic and dormant behavior enables identification and functional analysis of candidate proteins affecting tumor cell dormancy and metastasis. Cultured cancer cells are easy to expand and fractionate and currently provide the best source for the analysis of metastasis-associated cell surface proteins in cancer cells. The drawback of these models is the lack of proper microenvironment, which has been shown to play a crucial role in metastasis. Therefore, expression results obtained from the cell line models need further validation in animal models and in clinical samples. In addition, mechanistic analyses are required for in depth understanding on how these molecules affect the metastatic process.

Various methods including density gradient centrifugation and numerous chemical labeling techniques have been described for the isolation and enrichment of the cell surface and/or plasma membrane proteins for proteomic analyses (Elschenbroich et al., 2010; Leth-Larsen et al., 2010; Cordwell and Thingholm, 2010). Due to their accessibility, cell surface proteins of intact cells can be tagged with a membrane-impermeable biotin on amino acid residues located in extracellular space, which allows exploitation of the extraordinarily stable and non-covalent interaction between avidin and biotin in isolation and detection of the biotinylated cell surface proteins. Several commercial chemical biotinylation reagents, which


vary in their biotin moiety, spacer and reactive moiety, have been developed (Elia, 2008).

Importantly, by using labeling methods, all proteins accessible for the labeling reagent e.g., ligands bound to their receptors are isolated and analyzed with downstream applications. When adherent cell cultures are used as starting material, ECM proteins and secreted proteins bound to their ligands or ECM can also be labeled and isolated. Finally, the isolated cell surface proteins can be quantified and identified by proteomics methods to reveal differentially expressed proteins.

## **2.4 Nucleophosmin; oncogene, tumor suppressor or both?**

Nucleophosmin (NPM, B23, numatrin, NO38, hereafter referred as NPM) is a ubiquitously expressed multifunctional nucleolar phosphoprotein involved in a complex network of biological activities related to both growth suppression and proliferation. Importantly, it seems that deregulation in NPM homeostasis is often observed in tumors. NPM expression and gene integrity are frequently altered in human cancers and it has been attributed both tumor suppressive and oncogenic functions (Grisendi et al., 2006).

The human *NPM1* gene maps to chromosome 5q35 and contains 12 exons. Alternative splicing of the *NPM1* transcript results into two isoforms, B23.1 and B23.2 (Wang et al., 1993). In the full length B23.1 exon 9 is spliced to exon 11. The coding sequence stops at exon 12 resulting in a protein containing 294 amino acid residues. The shorter B23.2 isoform consists of 259 amino acids and lacks 35 amino acids at the C-terminus as a result of splicing the exon 9 to exon 10, which contains a stop codon. B23.2 is present in cells at low levels (Wang et al., 1993). Circular dichroism spectral analysis revealed similar secondary structures (mainly of beta-sheet and beta-turns) in B23.1 and B23.2 (Umekawa et al., 1993). Recently a third splice variant that lacks an internal exon 8 has been identified in the human EST database (GenBank accession number: NM\_199185).



**Figure 2.** Functional motifs and domains of NPM/B23. NPM is composed of a non-polar N-terminal domain containing nuclear export signal (NES). This part of the protein is important for oligomerization and molecular chaperone function. The central part of the protein contains a two acidic clusters required for histone binding. The C-terminal domain is needed for nucleic acid binding and together with the central region possesses ribonuclease activity. Adapted and modified from: (Grisendi et al., 2006).

The functional motifs and domains of NPM are shown in Figure 2. NPM contains motifs for nucleolar localization (Nishimura et al., 2002), nuclear import (Hingorani et al., 2000) as well as for nuclear export (Yu et al., 2006; Wang et al., 2005). It has distinct N- and C-terminal domains and a central part containing two acidic clusters important for NPM's histone binding (Okuwaki et al., 2001) and ribonuclease activity (Hingorani et al., 2000). Importantly, the C-terminal domain of NPM interacts with multiple proteins, like p53 (residues 249–262 of NPM) (Lambert and Buckle, 2006), c-Myc (residues 187–259 of NPM) (Li et al., 2008), FOXM1 (residues 187–259 of NPM) (Bhat et al., 2011) and Akt (residues 239–294 of NPM) (Lee et al., 2008). In addition, the C-terminal domain (residues 260–294 together with Thr199 and Thr234/237) is shown to be responsible for the binding of NPM to the phosphorylated RB (Takemura et al., 1999; Lin et al., 2010). Residues in parenthesis refer to the different fragments of NPM used to determine the binding site. Therefore, the actual binding site might be shorter. In addition,

NPM has been reported to bind to a number of other proteins involved in cellular processes like DNA replication, transcription, and repair, cell cycle control, ribosome biogenesis, viral replication, apoptosis, stability and splicing of mRNA, protein modification, mitotic spindle, cytoskeleton, and centromeres.

Notably, most of the functional data on NPM is supported by biochemical and *in vitro* data only. A future challenge is to understand the mechanisms which control NPM activation and recruitment to distinct subcellular sites and protein complexes to carry out its pleiotropic, often seemingly antagonistic, biological functions. The emerging picture indicates that NPM's biological roles are tightly regulated by several mechanisms such as NPM's expression level, localization, oligomerization status, post-translational modifications and NPM binding partners. Cellular distribution of NPM is closely associated with phospho/dephosphorylation events (Yun et al., 2003) and the nucleolar localization of NPM has been reported to require adenosine-5'-triphosphate (ATP) (Chang et al., 1998; Choi et al., 2008; Choi et al., 2008).

NPM undergoes several post-translational modifications, which regulate its cellular localization and function. A recent *in silico* analysis predicted that contains 40 potential phosphosites which would be substrates of at least 38 kinases. Based on the associated kinases the authors suggested that NPM phosphorylation is related to cellular processes such as apoptosis, cell survival, cell proliferation, and response to DNA damage stimulus (Ramos-Echazabal et al., 2012), all functions previously attributed to NPM. Post-translational modifications of NPM are summarized in table 1.

The following paragraphs summarize the current knowledge on NPM regulation and functions relevant to this thesis work.

**Table 1.** Post-translational modifications of NPM

<b>Enzyme</b>	<b>Modification</b>	<b>Timing</b>	<b>Relevance</b>	<b>Ref.</b>
<b>Plk2</b>	phosphorylation (serine 4)	S-phase	Centriole duplication	(Krause and Hoffmann, 2010)
<b>GRK5</b>	phosphorylation (serine 4)		sensitivity of cells to PLK1 inhibition	(So et al., 2012)
<b>Unknown kinase</b>	phosphorylation (serines 10 and 70)		Regulation of CDK1 activity	(Du et al., 2010)
<b>CK2</b>	phosphorylation (serine 125)	interphase	Dissociation from nucleolus	(Negi and Olson, 2006)
<b>ATR</b>	phosphorylation (serine 125)		Inhibition of the UV-induced p53 phosphorylation at Ser15	(Maugel et al., 2004)
<b>Cyclin B-CDK1</b>	phosphorylation (threonines 199, 219, 234, 237)	mitosis	Dissociation from the nucleolus, RNA binding, centrosomal association	(Peter et al., 1990; Okuwaki et al., 2002; Negi and Olson, 2006; Cha et al., 2004)
<b>Cyclin E-CDK2</b>	phosphorylation (threonine 199)	G1	Centrosome cycle control	(Okuda et al., 2000; Tokuyama et al., 2001)
<b>Cyclin A-CDK2</b>	phosphorylation (threonine 199)	S-phase and G1	Centrosome cycle control	(Tokuyama et al., 2001)
<b>Cyclin D-CDK4</b>	phosphorylation (threonine 199)	G1	Centrosome cycle control	(Adon et al., 2010)
<b>Viral cyclin V-CDK6</b>	phosphorylation (threonine 199)		Kaposi,s sarcoma herpesvirus latency	(Sarek et al., 2010)

Enzyme	Modification	Timing	Relevance	Ref.
<b>PKC</b>	phosphorylation (serine 225)		Phosphorylation <i>in vitro</i> in nuclear membrane fraction	(Beckmann et al., 1992)
<b>Nek2A</b>	phosphorylation	mitosis	Centrosomal localization	(Yao et al., 2004)
<b>PP1<math>\beta</math></b>	Dephosphorylation (threonines 199, 234, 237)		pRB binding and consequent E2F1-dependent DNA repair	(Negi and Olson, 2006; Lin et al., 2010; Haneji et al., 2012)
<b>SUMO-1 and SUMO-2</b>	Sumoylation		NPM localization, ribosome biogenesis	(Tago et al., 2005; Liu et al., 2007; Haindl et al., 2008)
<b>SENP3 and 5</b>	Desumoylation		Control of ribosome biogenesis	(Haindl et al., 2008; Yun et al., 2008)
<b>p300</b>	Acetylation mostly on the C-terminal domain)		Stimulation of chromatin transcription	(Swaminathan et al., 2005)
<b>BRCA1-BARD1 ubiquitin ligase</b>	ubiquitinylation	mitosis	Stabilization of NPM	(Sato et al., 2004)
<b>Not reported</b>	poly-(ADP-ribosyl)ation			(Ramsamooj et al., 1995)
<b>Plk2 = Polo-like kinase, GRK5 = G protein-coupled receptor kinase 5, CK2 = Casein kinase 2, ATR = Ataxia telangiectasia mutated and Rad3 Related protein kinase, PKC = Protein kinase C, PP1 <math>\beta</math> = Protein phosphatase 1 <math>\beta</math>, SUMO = Small ubiquitin-like modifier, SENP = SUMO1/sentrin/SMT3 specific peptidase</b>				

#### **2.4.1 Discovery of NPM as nucleolar protein involved in ribosome assembly and transport**

NPM was originally designated as B23 based on its position on a two dimensional gel analysis from acid-extracted nucleolar proteins isolated from normal rat liver and Novikoff hepatoma ascites cells (Orrick et al., 1973). It was shown to be a highly expressed nucleolar phosphoprotein (Kang et al., 1974; Prestayko et al., 1974), which localizes especially to the granular region of the nucleolus, a site where ribosomes are assembled (Spector et al., 1984). In addition, NPM was proved to be identical with a nuclear protein numatrin, which is tightly bound to nuclear matrix (Feuerstein and Mond, 1987; Feuerstein et al., 1988). NPM's nucleolar localization and its shuttling between the nucleus and cytoplasm (Borer et al., 1989) led to the earliest proposal that it facilitates ribosome assembly (Dumbar et al., 1989) and transport of pre-ribosomal particles (Yung et al., 1985). Later NPM has been shown to direct the nuclear export of both 40S and 60S ribosomal subunits (Maggi et al., 2008). According to current knowledge, NPM provides the necessary export signals and chaperoning capabilities that are required to transport ribosome components from nucleus to cytoplasm and by these functions it seems to balance protein synthesis to cell growth and proliferation (Grisendi et al., 2006; Falini et al., 2007), important determinants of cancer growth.

#### **2.4.2 NPM as a chaperone and its role in transcriptional regulation**

Genomic DNA is compacted into chromatin by organizing it into nucleosomes by its association with four histone proteins H2A, H2B, H3, and H4. The correct assembly/disassembly of nucleosomes is mediated by histone chaperones whose precise function is necessary for DNA-dependent activities like transcription, replication and repair (Burgess and Zhang, 2013).

NPM is biochemically defined as a member of the nucleoplasmin/nucleophosmin family of nuclear chaperones. The family

members share an acidic core-domain at the N-terminus of the protein which is responsible for oligomer formation and chaperone activity (Schmidt-Zachmann et al., 1987; Hingorani et al., 2000; Frehlick et al., 2007; Okuwaki et al., 2001; Prinos et al., 2011). Crystal structure of the N-terminal core domain in NPM revealed that it forms pentamers which can further oligomerize into decamers similar to the other nucleoplasmin family members (Lee et al., 2007). In addition to the N-terminal domain, some C-terminal regions might be required for NPM oligomer formation (Liu and Chan, 1991). A recent computational analysis of the N-terminal domain in NPM proposes that NPM monomer-oligomer status can be regulated by transformation from a folded, pentameric structure to a monomeric, disordered state through phosphorylation events (Mitrea and Kriwacki, 2012).

As a nuclear chaperone NPM can bind to histones and assemble nucleosomes *in vitro* (Okuwaki et al., 2001; Gadad et al., 2011). NPM possesses chaperone activity also for proteins since it can prevent aggregation and thermal denaturation of proteins such as HIV-1 Rev protein, liver alcohol dehydrogenase and carboxypeptidase A *in vitro* (Szebeni and Olson, 1999).

NPM's histone chaperone activity is enhanced by p300 mediated acetylation on the C-terminal domain which modulates *in vitro* transcription from chromatin templates by RNA Pol II (Swaminathan et al., 2005). Furthermore, acetylated NPM localizes to nucleoplasm where it regulates transcriptional activation of genes implicated in oral cancer manifestation such as tumor necrosis factor alpha and interleukin-6 receptor (Shandilya et al., 2009). NPM negatively regulates a histone-modifying enzyme GCN5 and thus transcription and this regulation was enhanced by phosphorylation of NPM at Thr199 (Zou et al., 2008). Recent results indicate that the NPM might regulate gene expression at specific G-quadruplex regions (Xu et al., 2007; Federici et al., 2010; Gallo et al., 2012), which are common in oncogene promoters, whereas a reduced frequency is observed in tumor suppressor genes (Qin and Hurley, 2008). NPM has been shown to regulate transcription, either positively or negatively, also through interaction with



several transcriptional regulatory partners like oncogenic transcription factor Forkhead box M1 (FOXM) (Bhat et al., 2011), androgen receptor (Leotoing et al., 2008), activating protein transcription factor 2 (Liu et al., 2007) and YY1 (Inouye and Seto, 1994), to mention some. Intriguingly, YY1 has also been indicated to regulate NPM gene expression (Chan et al., 1997).

NPM associates with both DNA and RNA with its C-terminal nucleic acid binding domain (Wang et al., 1994; Hingorani et al., 2000; Okuwaki et al., 2002) and it has been reported to have endoribonuclease activity to ribosomal RNA (rRNA) (Savkur and Olson, 1998; Hingorani et al., 2000). NPM binds to rRNA chromatin and regulates the histone density around rRNA genes (Murano et al., 2008). This activity requires NPM's RNA binding activity and is regulated by its cell cycle-dependent phosphorylation (Hisaoka et al., 2010).

The B23.2 splice variant of NPM lacks the C-terminal domain and therefore is not able to bind double stranded DNA (Wang et al., 1994) and has lower ribonuclease activity than the full-length NPM protein (Herrera et al., 1995). In addition, B23.2 can heterodimerize with the full length NPM and reduces its RNA-binding (Okuwaki et al., 2002). Oligomer formation by itself has also been shown to decrease NPM's DNA binding (Herrera et al., 1996).

#### **2.4.3 Molecular pathways regulated by NPM: RB, ARF-p53-Mdm2 pathway and c-Myc**

The retinoblastoma protein (RB), encoded by the *RB1* gene, is the first known human tumor suppressor (Knudson, 1984; Corson, 2007; Dimaras, 2008; Huang et al., 1988) which is inactivated in multitude of solid cancers by various distinct mechanisms (Burkhart and Sage, 2008). In normal quiescent tissues RB maintains cell cycle arrest by repressing the activity of E2F-family of transcription factors (E2F1, E2F2, and E2F3) (Burkhart and Sage, 2008; Cobrinik, 2005). This repression is relieved by mitogenic or oncogenic signals, which induce RB phosphorylation (pRB) (Mittnacht, 1998). As a

consequence, the expression of genes involved in DNA-replication, mitosis and cytokinesis is induced (Markey et al., 2007).

Hyperphosphorylated RB translocates into nucleoli in late S or G<sub>2</sub> phase. Translocation is mediated by NPM (Takemura et al., 1999; Takemura et al., 2002), most likely after dephosphorylation of NPM on threonines 199, 234, 237 by PP1 $\beta$  (Lin et al., 2010). The biological significance of the pRB-NPM complex and nucleolar localization is unclear but RB and NPM have been shown to synergistically stimulate DNA polymerase alpha activity (Takemura et al., 1999).


The p53 tumor suppressor gene is defective in about half of all tumors, regardless of their type or origin (Hollstein et al., 1991; Levine et al., 1991) and in the remaining cancers a considerable number has alterations in the p53 pathway. The p53 gene encodes a tetrameric protein that functions mainly as a transcription factor at the crossroads of cellular stress response pathways (like DNA damage, oncogene activation and hypoxia) controlling the expression of genes involved in cell division and viability, growth arrest and apoptosis (Levine and Oren, 2009). NPM regulates p53 by directly binding to it (Colombo et al., 2002; Maignel et al., 2004) and by affecting the p53 regulatory proteins ARF (Bertwistle et al., 2004; Korgaonkar et al., 2005) and Mdm2 (Kurki et al., 2004). NPM has a dual effect on p53 and it can either stabilize (Colombo et al., 2002; Horn and Vousden, 2004; Kurki et al., 2004) or inhibit p53 (Li et al., 2004; Wu et al., 2002; Dhar and St Clair, 2009; Li et al., 2007) and apoptosis. Mechanisms regulating such opposite effects are mainly unknown. NPM oligomerization might be necessary for its capability to inhibition p53-mediated apoptosis (Qi et al., 2008; Jian et al., 2009). NPM could also inhibit p53 activity by competing with p53 phosphorylation (Maignel et al., 2004; Nalabothula et al., 2010). NPM binding partners have also been reported to regulate its effect on p53 (Ji et al., 2012; Fukawa et al., 2012).

The small tumor suppressor protein p14ARF (humans, p19ARF in mouse, hereafter referred as ARF) functions as a tumor suppressor by inhibiting the Mdm2 mediated degradation of p53. It binds directly to Mdm2 in a site distinct from the p53 binding domain (Ashcroft and Vousden, 1999) which

inhibits the ubiquitin ligase activity of Mdm2 (Honda and Yasuda, 1999). Even though the complex interaction between ARF and NPM has been under extensive investigation over the last years, it is still not fully understood. NPM binds ARF in the nucleoli in a quantitative manner (Bertwistle et al., 2004). ARF mutants lacking the NPM binding site are unstable (Kuo et al., 2004) and ARF localizes to nucleoplasm in MEFs lacking both p53 and NPM (Colombo et al., 2005) indicating that NPM stabilizes ARF in the nucleoli. On the other hand, ARF has been shown to mediate NPM degradation (Itahana et al., 2003) indicating a feed-back loop between these two proteins. ARF has also been shown to block NPM function in rRNA processing and the transport of pre-ribosomal particles (Savkur and Olson, 1998; Itahana et al., 2003; Sugimoto et al., 2003; Brady et al., 2004). NPM is also a target for ARF induced sumoylation (Tago et al., 2005).

c-Myc induces both p53-dependent and p53-independent apoptosis upon up-regulation of the tumor suppressor ARF. However, when overexpressed or deregulated c-Myc becomes oncogenic (Hoffman and Liebermann, 2008; Nilsson and Cleveland, 2003). Importantly, in animal models, most if not all, c-Myc-induced tumors have inactivated the ARF-p53 pathway (Eischen et al., 1999; Nilsson and Cleveland, 2003). There is a complex interaction between NPM and the c-Myc-oncogene. Expression of c-Myc correlates with NPM expression (Guo et al., 2000; Neiman et al., 2001; Kim et al., 2000) and NPM is a transcriptional target of c-Myc (Zeller et al., 2001). In addition, NPM directly interacts with c-Myc and regulates c-Myc mediated rDNA transcription, nucleolar localization (Li and Hann, 2013) and expression of c-Myc target genes like eIF4E (Li et al., 2008). NPM overexpression has been shown to enhance c-Myc-induced proliferation and transformation in p53<sup>-/-</sup> ARF<sup>-/-</sup> double knockout (DKO) mouse embryo fibroblasts (MEFs) (Li et al., 2008)

Recent studies, however, have shown that NPM binds to the G-quadruplex DNA on c-Myc gene-promoter which suppresses c-Myc gene expression (Siddiqui-Jain et al., 2002; Gallo et al., 2012). In addition, Reduction in NPM levels accelerated lymphomagenesis in  $\mu$ -Myc transgenic mice (Grisendi et al., 2005).



**Figure 3.** Schematic representation of the discussed pathways controlled by NPM. Solid lines indicate connections with strong scientific background.

#### 2.4.4 Role of NPM in centrosome duplication

Centrosomes, the microtubule-organizing centers of animal cells, are mainly composed of mother and daughter cylindrical microtubule-based centrioles and a matrix of associated pericentriolar material. Centrosome is duplicated in coordination with DNA replication. Centrosome amplification is frequently observed in cancers and is postulated to be one cause of chromosome instability. Proper centrosome function is also required for cytokinesis (Wang et al., 2004; Zyss and Gergely, 2009).

Cultured NPM-null cells or mice carrying a single inactivated *Npm1* allele display increased centrosome numbers (Grisendi et al., 2005). In accordance, NPM is shown to control centrosome cycle. NPM attaches to single, unduplicated centrosomes in late G1 (Ochs et al., 1983) and re-associates with the centrosomes during mitosis (Zatsepina et al., 1999; Okuda et al., 2000) at least partly after its phosphorylation on Thr234 and Thr237 by cyclin B-CDK1 complex (Cha et al., 2004). At the centrosome, NPM localizes between the centriole pair (Shinmura et al., 2005). Cyclin E(A)-CDK2 (Tokuyama et al., 2001; Okuda et al., 2000) and cyclin D-CDK4 phosphorylate NPM at Thr199, which partly dissociates NPM from the

centrosome subsequently allowing separation of the centrioles and centrosome duplication (Okuda et al., 2000; Tokuyama et al., 2001; Okuda, 2002; Tarapore et al., 2002).

Some NPM phosphorylated on Thr199 remain attached to centrioles, translocates toward a mother centriole (Shinmura et al., 2005) and binds to and activates centrosomal ROCK II kinase, which further promotes centrosome duplication (Ma et al., 2006). In addition to Thr199 phosphorylation, Plk2-mediated phosphorylation of NPM on serine 4 has been shown to promote centrosome duplication (Krause and Hoffmann, 2010). However, while important in regulating the correct centrosome cycle, the Thr199 phosphorylation might not be necessary for NPM's growth promoting functions (Brady et al., 2009).

#### **2.4.5 NPM in human tumors and in mice**

NPM protein levels are elevated in rapidly dividing cells (Feuerstein and Mond, 1987; Feuerstein and Mond, 1987; Nozawa et al., 1996) and its expression increases rapidly in the early G1 phase during mitosis (Feuerstein et al., 1988) which led to the proposed growth promoting function of NPM. In accordance, NPM levels decrease in cells that are withdrawn from the cell cycle or undergoing apoptosis (Hsu and Yung, 1998; You et al., 1999; Yung, 2004). NPM expression associates with poor prognosis, recurrence or lymph node metastasis in Ewing's sarcoma (Kikuta et al., 2009), bladder carcinoma (Tsui et al., 2008), colon cancer (Liu et al., 2012) and oral squamous cell carcinoma (Coutinho-Camillo et al., 2010) in a relative small patient materials (n=31-154). In hematological malignancies NPM is involved in chromosome translocations, where its N-terminal oligomerization domain is fused with different partners like ALK, RAR $\alpha$  and MLF. NPM oligomerization domain contributes to tumor development by activating the oncogenic potential of the fused protein partner (Grisendi et al., 2006).

On the contrary, Bocker and coworkers did not find difference in NPM expression between normal and malignant epithelial prostatic cells (Bocker

et al., 1995) and NPM expression did not correlate with patient prognosis in an analysis of 55 melanoma patient samples (Calli et al., 2011). No amplification of *NPM1* has been detected in human tumors, but deletion of the gene has been observed in non-small cell lung carcinomas (Mendes-da-Silva et al., 2000). In addition, *NPM1* is the most frequently mutated gene in acute myeloid leukemia (AML) accounting for ~35% of cases (Falini et al., 2005). Mutations in the C-terminal domain lead to mutant proteins that aberrantly and stably localize to the cytoplasm due to an additional nuclear export signal at the C-terminal domain (Bolli et al., 2007; Albiero et al., 2007). Mutant NPM oligomerizes with the wild type protein, which is also transported to the cytoplasm thus functioning as a dominant negative (Falini et al., 2009). These mutations seem to represent tumor –initiating lesions, since they appear before the other AML-associated genetic alterations can be detected (den Besten et al., 2005).

*Npm1* acts as a haploinsufficient tumor suppressor in the hematopoietic compartment in mice (Sportoletti et al., 2008). Inactivation of *Npm1* in the germ line leads to embryonic lethality in mice at midgestation indicating indispensable role during development (Colombo et al., 2005; Grisendi et al., 2005). *Npm1* null embryos display genomic instability, widespread apoptosis, and activation of p53. In addition, cultured NPM-null cells or mice carrying a single inactivated *Npm1* allele accumulate DNA damage (Colombo et al., 2005) and have increased centrosome numbers (Grisendi et al., 2005). In addition, *Npm1*<sup>+/-</sup> MEFs have an immortal phenotype with noticeably high proliferation rates (Colombo et al., 2005; Grisendi et al., 2005) and display accelerated lymphomagenesis when crossed with  $\mu$ -Myc transgenic mice (Grisendi et al., 2005).


### 3 AIMS OF THE STUDY

Cell surface proteins are known to be important players in regulating cancer metastasis. As the molecular mechanisms underlying the second phase of metastasis (colonization at the distant site) and metastatic dormancy are poorly resolved, we wanted to identify novel cell surface proteins contributing to colonization at the distant site. The aim was not to gain a comprehensive picture of the cell surface proteins of the two cell lines used but to identify differentially expressed proteins for future studies.

The specific aims included:

- Isolate the cell surface proteins from invasive, non-metastatic and fully metastatic cell clones of the MDA-MB-435 cell line for a proteomic based comparative study
- Profile the differential expression of cell surface proteins in order to discover interesting, differentially expressed proteins for future functional studies in metastatic colonization
- Analyze the breast cancer associated role and function of one of the identified proteins, nucleophosmin, in more detail

## 4 MATERIALS AND METHODS



**Figure 4.** Flowchart of the study



The materials and methods of this study are listed in table 2. with detailed descriptions in the following text and in the original publications, here referred to by Roman numerals.

**Table 2.** Methods used in the study

<b>Methods</b>	<b>Used in</b>
<b>Cell proliferation assays</b>	II, III
<b>Cell surface biotinylation</b>	I, unpublished data
<b>Cloning</b>	II, III
<b>Creation of stable cell lines</b>	II, III
<b>Gene knockdown by RNA interference</b>	III
<b>Immunofluorescence microscopy</b>	I, II, III, unpublished data
<b>Immunohistochemistry</b>	I, II, III
<b>Immunoprecipitation</b>	II, III
<b>Implantation of tumor cells into mice</b>	I
<b>Lentiviral production</b>	II
<b>Mammalian cell culture</b>	I, II, III
<b>Mass spectrometry based protein identification</b>	I, unpublished data
<b>Network and pathway analyses</b>	II
<b>Quantitative real-time PCR</b>	II
<b>RNA extraction</b>	II, III

Methods	Used in
Scoring of tumor microarrays	II, III
SDS-PAGE and immunoblotting	I, II, II, unpublished data
Sequence analysis	II, III
Soft agar assay	II
Statistics	I, II, III
Transduction of mammalian cells	II
Transfection of mammalian cells	II, III
3D cell culture	II
17 $\beta$ -Estradiol Treatment	II

#### 4.1 Cell culture (2D and 3D), cell treatments, cell proliferation and soft agar assays

Human cancer cell lines (Table 1) were cultured at + 37 °C in 5% CO<sub>2</sub> atmosphere in appropriate medium. For 3D growth/invasion assays cross-linked fibrin gels were prepared by combining 75  $\mu$ l plasminogen-free human fibrinogen (6 mg/ml; Calbiochem) in Hank's Balanced Salt Solution (HBSS) and 75  $\mu$ l HBSS (pH 7.4) containing 4 U/ml human thrombin and 400  $\mu$ g/ml aprotinin (both from Sigma-Aldrich). 5000 cells were suspended in 40  $\mu$ l of the prepared, cross-linked fibrin and transferred to a 24-well plate. The plate was incubated for 1 h at 37°C to allow complete gelling. The cells were grown for 4–16 days, fixed and photographed

For the 17 $\beta$ -Estradiol Treatment cells were starved for 48 hours in Phenol Red-free RPMI (Gibco, California, USA) supplemented with 2% charcoal-dextran filtrated fetal calf serum (HyClone; Thermo Scientific, Logan, UT), 1% glutamine, and penicillin-streptomycin. Starved cells were treated with 10 nmol/L 17 $\beta$ -estradiol for 3 days and analyzed.

Cell proliferation was measured by growing cells on 96-well plates for indicated time periods. Cells were incubated 2 hours with 10  $\mu$ L MTT (5 mg/mL) and lysed (10% SDS, 10 mmol/L HCl) overnight. Absorbance was measured at 540 nm using Multiskan Ascent software version 2.6 (Thermo Labsystems, Vantaa, Finland).

For Soft agar assays tissue culture dishes containing a 2-mL layer of solidified 0.7% agar in a complete medium were prepared. Cells ( $5 \times 10^3$  cells per 35-mm well) were suspended in complete medium containing 0.35% agarose and plated on top of the solidified bottom agar. After 14 days, number of colonies was quantified. For visualization, foci were methanol-fixed and stained with 0.005% crystal violet.

## **4.2 Cell surface protein isolation**

Cells were grown 3 days after which they were washed three times with Dulbecco (PBS + 0.901 mM  $\text{CaCl}_2$  + 0.492 mM  $\text{MgCl}_2$ ) and labeled with EZ-Link Sulfo-NHS-SS-Biotin (Pierce, Rockford, IL, USA; 0.5 mg/ml in) for 30-minutes in Dulbecco on ice. Labeling solution was removed by washing the cells twice with Dulbecco followed by blockade of the non-reacted biotin with 20 mM glycine for 15 minutes and washed with Dulbecco + 100  $\mu$ M oxidized glutathione (Sigma-Aldrich, St. Louis, MO). Cells were lysed and the membrane proteins solubilized in 500  $\mu$ L of lysis buffer (PBS + 2% NP-40, 1% Triton X-100, 10% glycerol, 100  $\mu$ M oxidized glutathione, EDTA free protease inhibitor tablet (Roche, Mannheim, Germany)) for 30 minutes. The cell extracts were incubated with 30 U of DNase (22  $^\circ\text{C}$  50 min, Roche, Mannheim, Germany) and centrifuged for 20 min (20 800  $\times g$ , at + 4  $^\circ\text{C}$ ). Equal amounts of protein from each cell extract, determined by DC- (Bio-Rad, Hercules, CA, USA) or BCA- (Thermo Fisher Scientific, Pierce Rockford, IL, USA) protein assays were used for cell surface protein isolation. The supernatant was pre-cleared using biotin agarose beads (ImmunoPure $^\circledR$  Immobilized D-biotin, Pierce, Rockford, IL, USA) and incubated with magnetic streptavidin beads (Dynabeads $^\circledR$ MyOne $^\text{TM}$  Streptavidin, Invitrogen Dynal, AS, Norway). Beads were washed four times with the lysis buffer, four times with 300 mM NaCl in lysis buffer and twice with 50 mM Tris-HCl, pH 7.8. and the isolated proteins were eluted with 50 mM DTT in 50 mM Tris-HCl, pH 7.8 at 30  $^\circ\text{C}$ , followed by pooling of the eluates.

### 4.3 Mass spectrometry, protein identification, network and pathway analyses

For LC–MS/MS analysis proteins were digested with trypsin (sequence grade modified, Promega, Madison, WI, USA) and peptides were loaded to a reversed phase pre-column (NanoEase Atlantis dC18, 180  $\mu$ m  $\times$  23.5 mm, Waters, Milford, MA, USA) with 0.1% formic acid followed by separation in reversed phase analytical column (PepMap 100, 75  $\mu$ m  $\times$  150 mm, Thermo Fisher Scientific Dionex, Germering, Germany) with a linear gradient (4–50%) of 95% acetonitrile in 0.08% formic acid in 35 min. Full scan for eluting peptides was acquired in an LTQ Orbitrap XL mass spectrometer (Thermo Fisher Scientific Inc.) introduced to via the ESI Chip interface (Advion BioSciences Inc.) in the positive-ion mode. The mass range of 300–2000 m/z on the Orbitrap-detector with 60 000 resolution (FWHM) at 400 m/z, the AGC target was set to 200 000 and the maximum inject time to 800 ms. Six MS/MS data-dependent scans were acquired on LTQ based on full scan, with the AGC target set to 10 000 and the maximum inject time set to 100 ms. An isolation width of 2 m/z was used for precursor selection. Peptide fragmentation was done in normalized collision energy of 35%, activation time of 100 ms and activation Q set to 0.25. Precursors, whose charge state couldn't be determined or charge state was + 1, were discarded from the MS/MS analysis. Precursors were dynamically excluded for 10 s with a repeat count of 1. Both full scan and MS/MS scans consisted of one microscan and they were acquired as profile data.

Mascot (version 2.2, in-house server), Swissprot 2010 (516 080 sequences, 181 677 051 residues) and the following parameters were used for protein identification: enzyme: trypsin, peptide mass tolerance: 0.02 Da, fragment mass tolerance 0.8 Da, max missed cleavages 2, variable modifications: carbamidomethyl (C), carboxymethyl (C), deamidated (NQ), oxidation (M), propionamide (C), and pyridylethyl (C). Same search was also conducted limiting the taxonomy to human, except using variable modifications carbamidomethyl (C), oxidation (M), and propionamide (C). Both results were combined and the results filtered to human proteins. Mascot score > 40 was required for identifications in the total protein list. In addition, at least one unique peptide was required for the reported differentially expressed proteins. Pathway and network analyses for the differentially expressed proteins were performed in a computational platform Moksiskaan.

## 4.4 Immunoprecipitation and immunoblotting

Equal amounts of protein from post nuclear supernatans of each extract were pre-cleared using protein G sepharose™ 4 fast flow beads (GE Healthcare Bio-Sciences Ab, Uppsala, Sweden) followed by incubation with the appropriate antibodies (+ 4 °C, o/n). The antibody-antigen complex was captured on G sepharose beads (+ 4 °C, 30 min), which were washed four times with lysis buffer and boiled in *Laemmli* buffer.

Proteins were separated using SDS-PAGE and transferred to Immobilon-P Membrane (Millipore, Bedford, MA, USA). Immunoblots were incubated with the appropriate primary and secondary antibodies (HRP conjugated, Dako, Glostrup, Denmark) followed by enhanced chemiluminescence visualization using the SuperSignal West Pico kit (Thermo Fisher Scientific, Pierce, Rockford, IL, USA).

## 4.5 Immunohistochemistry and immunofluorescence analyses

For immunofluorescence staining of intracellular antigens, cells were fixed with 4% paraformaldehyde and permeabilized with 0.5% NP-40 in PBS. Unspecific antibody binding sites were blocked using 3% bovine serum albumin in PBS. The cells were incubated with primary and secondary (Molecular Probes goat anti-mouse Alexa Fluor 594; Invitrogen, Carlsbad, CA) antibodies. For staining of cell surface antigens live cells were blocked and incubated with the primary antibodies on ice after which they were fixed and incubated with the secondary antibodies. DNA was visualized with DAPI (Vector Laboratories, Burlingame, CA).

For the immunohistochemical analyses paraffin embedded sections were deparaffinized and the antigens retrieved with the appropriate method for each antibody. Primary antibodies were detected by using the TSA-kit (Perkin Elmer, Waltham, MA, USA) and the signal was visualized with the AEC-reagent (Sigma-Aldrich, St. Louis, MO) or by using peroxidase staining (biotinylated horse anti-mouse immunoglobulins 1:200 and Vectastain ABC complex; Vector Laboratories), which was visualized with 3,3'-diaminobenzidine (Vector Laboratories). Images were captured with an Olympus DP50 camera and with Olympus Studio Lite software version 1.0 or 1.01. For the C-kit staining we used PowerVision Novocastra preantibody blocking solution (Leica Microsystems, Wetzlar, Germany) for primary antibody incubation. Staining was detected using a PowerVision + Poly-HRP histostaining kit (DPVB + 110DAB; ImmunoVision Technologies, Daly City,

CA; Springdale, AR) according to the manufacturer's instructions. A Ventana Discovery IHC slide stainer (Ventana Medical Systems, Tucson, AZ) and a Ventana 3,3'-diaminobenzidine tetrahydrochloride biotin avidin detection kit was used for the cytokeratin staining.

#### **4.6 Patients and classification of breast cancer subtypes in the TMAs**

We used two different breast cancer patient cohorts, the FinProg and FinHer, which both contained detailed clinical information allowing survival and association analyses. The cancers were classified into five biological subtypes as follows; luminal A (ER+ and/or PR+, HER2-), luminal B (ER+ and/or PR+, HER2+), basal-like (ER-, PR-, HER2-, CK5+ and/or EGFR+), HER2-enriched (HER2+, ER-, PR-), and non-expressor type (negative for all five key classifiers). The non-expressor type corresponds to unclassifiable triple-negative breast cancer. An ethics committee at Helsinki University hospital approved the FinHer study. Regarding FinProg material permission to use formalin fixed paraffin embedded tissues for research purposes was provided by the Ministry of Social Affairs and Health, Finland.

#### **4.7 Statistical analyses**

In the TMAs the associations between factors were tested with the  $\chi^2$  test. The odds ratio was used to examine the strength of the relationships. Kaplan-Meier method was applied to calculate the life tables. Multivariate survival analyses were performed with the Cox proportional hazards model, entering the appropriate covariates.

Progenesis LC-MS software was used to quantitate the expression differences in the comparative proteomic analysis. The runs were divided into following 4 groups; metastatic and non-metastatic non-biotinylated controls and metastatic and non-metastatic samples. The peptides present in control groups, determined by manual comparison of the peptides, were discarded from the analysis followed by comparison of the metastatic and non-metastatic samples against each other. ANOVA and q-values calculated by the Progenesis software were used to deduce differentiating peptides. The Progenesis stats-package was used to perform unsupervised principal component analysis (PCA).

## **5 RESULTS AND DISCUSSION**

### **5.1 Comparative analysis of metastasis and dormancy associated cell surface proteins (I and unpublished data)**

As described earlier, the details and molecular mechanisms of metastasis are not fully resolved. Therefore, we wanted to identify novel cell surface proteins contributing to the formation of metastasis, especially to the second phase (colonization at the distant site). In this thesis work we used metastatic and non-metastatic subclones of the MDA-MB-435 human carcinoma cell line as a model for the metastatic spread of cancer. The non-metastatic cells leave the primary tumor as efficiently as the metastatic ones and are detected as single cells in the lung capillaries. However, they fail to go through the latest steps of the metastatic cascade and cannot form full-blown metastases (Goodison et al., 2003).

Originally, this thesis work aimed at comparing the differences in cell surface proteins between the metastatic and non-metastatic cell pair by using two-dimensional (2D) gel electrophoresis. (Karhemo et al., unpublished data). However, the fast development of gel-free shotgun proteomics technologies (Duncan et al., 2010) and mass spectrometry based protein quantification (Ong and Mann, 2005; Domon and Aebersold, 2010) led us to analyze and quantify our isolated cell surface samples using liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS) (Study I).

### **5.1.1 Optimized method for isolation of biotinylated cell surface proteins (I)**

For comparative proteomic analysis the cell surface proteins of the isogenic non-metastatic and metastatic cell pair were labeled with biotinylation reagents that react with primary amines (-NH<sub>2</sub>), such as the side-chain of lysine followed by isolation with magnetic streptavidin beads.

Studies used as reference to optimize the isolation of biotinylated cell surface proteins reported a consistent identification of intracellular proteins (Scheurer et al., 2005; Roesli et al., 2008). To diminish this intracellular background the cell extracts were treated with DNaseI to dissolve the cytosolic actin and proteins associated with it from the viscous DNA in order to liberate a maximal amount of membrane proteins. In addition, samples were pre-cleared by using biotin agarose beads and intracellular background was controlled by including non-biotinylated control samples in the analysis.

The gel-free LC-MS/MS approach resulted in identification of 86 proteins in total from both cell lines. This relatively limited number of identifications might be explained by performance of database search within the Progenesis LC-MS software after exclusion of all peptides present in the control samples, by the low amount of starting material and the performance of the LC-MS analyses without exclusion lists to exclude already fragmented features.

Importantly, more than 60% of the identified proteins localized to the cell surface/extracellular space according to the Finnish Red Cross in house Cell Surface Protein Classifier. The specific enrichment of cell surface proteins was further verified by analyzing the presence of a cell surface protein, cluster of differentiation 109 (CD109), the endoplasmic protein BiP, the Golgi protein Golgin 97, the subunit IV of mitochondrial protein cytochrome C oxidase and an intracellular protein Erk1/2 in the cell surface and unbound fractions (representing proteins that did not bind to the streptavidin beads) in a Western blot analysis. Our optimized protocol specifically isolated the cell surface proteins since that fraction was free of organelle contaminants (Figure 1A and B in publication I).



### **5.1.2 Cell surface protein changes in the metastatic cells (I, unpublished data)**

This thesis work revealed 29 differentially expressed proteins (Table 3) from the cell surface samples between the metastatic and non-metastatic cells using the 2D gels (Karhemo et al, unpublished results) and the LC-MS/MS analyses (publication I). Six differentially expressed proteins were identified from silver stained 2D gels based on visual comparison of the spot intensities (Table 3). The gel-free analysis revealed 23 differentially expressed proteins (Table 3, Table 1 in publication I).

Based on a Uniprot database (<http://www.uniprot.org/>) annotation and literature search 19/29 proteins (66%) represented proteins with known cell surface localization. The gel-free LC-MS/MS based approach proved to be superior in identifying cell surface membrane proteins compared to the 2D gels. Only one protein, LGALS3, identified using the 2D gels is reported to be secreted to the cell surface and only one protein is predicted to contain a single transmembrane domain. On the contrary, 18/23 (78%) of the differentially expressed proteins identified in the gel-free analysis represented proteins with previously reported cell surface localization and 15 were either single or multi-pass transmembrane proteins (Table 3).

Interestingly, most of the identified proteins (20/29, 69%) were overexpressed in the metastatic cells. Similar results have been obtained in a previous proteomic analysis of ultracentrifuged and density separated plasma membrane fractions of the same isogenic MDA-MB-435 cancer cell pair (Lund et al., 2009). These results indicate that overexpression/localization of certain proteins on the surface of the metastatic cells may facilitate extravasation and colonization of the metastatic cells by adapting them to the new microenvironment at the secondary site.

**Table 3.** Differentially expressed proteins identified in the thesis work

Name	Relative expression (metastatic/non-metastatic) in LC-MS/MS	Method	Cellular localization <sup>A</sup>	Type of membrane protein
<b>MRPL12</b>		2D	mitochondrion, ribosome	
<b>RPLPo</b>		2D	cytoplasm, ribosome	
<b>MRPL39</b>		2D	mitochondrion, ribosome	
<b>NPM</b>		2D	nucleolus, nucleus	
<b>SSR4</b>		2D	endoplasmic reticulum membrane	single-pass
<b>LGALS3</b>		2D	secreted, cytoplasm, nucleus	peripheral
<b><u>HLA-DRB1</u></b>	8.8	LC-MS/MS	cell membrane	single-pass
<b><u>HLA-DRA1</u></b>	2.53	LC-MS/MS	cell membrane	single-pass
<b><u>BST2</u></b>	7.88	LC-MS/MS	cell membrane, trans-Golgi network	single-pass, GPI-anchor
<b><u>PTGFRN</u></b>	7.44	LC-MS/MS	cell membrane	single-pass
<b><u>ART3</u></b>	6.03	LC-MS/MS	cell membrane	GPI-anchor
<b><u>PROCR</u></b>	4.4	LC-MS/MS	cell membrane	single-pass
<b><u>GPR56</u></b>	3.01	LC-MS/MS	cell membrane	multi-pass
<b><u>ADAM10</u></b>	2.74	LC-MS/MS	cell membrane	single-pass
<b><u>PTPRF</u></b>	2.4	LC-MS/MS	cell membrane	single-pass
<b><u>ATP1B3</u></b>	2.21	LC-MS/MS	cell membrane	single-pass
<b><u>CD109</u></b>	2.11	LC-MS/MS	cell membrane	GPI-anchor


Name	Relative expression (metastatic/non-metastatic) in LC-MS/MS	Method	Cellular localization <sup>A</sup>	Type of membrane protein
<b><u>ITGA6</u></b>	2.08	LC-MS/MS	cell membrane	single-pass
<b><u>TUBA1C</u></b>	2.01	LC-MS/MS	cytoplasm, cytoskeleton	
<b><u>IGSF8</u></b>	2	LC-MS/MS	cell membrane	single-pass
<b>ITGAV</b>	1.82	LC-MS/MS	cell membrane	single-pass
<b>HSPD1</b>	1.76	LC-MS/MS	mitochondrion matrix	
<b>FASN</b>	1.7	LC-MS/MS	cytoplasm	
<b>ITGB1</b>	1.44	LC-MS/MS	cell membrane	single-pass
<b><u>GOT2</u></b>	0.43	LC-MS/MS	mitochondrion matrix, cell membrane	peripheral
<b><u>NME1</u></b>	0.49	LC-MS/MS	cytoplasm, nucleus	
<b>CHL1</b>	0.56	LC-MS/MS	cell membrane	single-pass
<b>CD97</b>	0.73	LC-MS/MS	cell membrane	multi-pass
<b>ALDH7A1</b>	0.75	LC-MS/MS	mitochondrion, nucleus	
<b>Overexpressed, underexpressed</b> (metastatic/non-metastatic)				
<sup>A</sup> UniProt and literature				
Proteins that displayed $\geq 2$ -fold in their expression are <u>underlined</u> , proteins whose differential surface expression was validated by antibodies are <b>highlighted</b> .				

Previous gene array analysis, which compared a subclone of the original metastatic MDA-MB-435 clone to the non-metastatic one, revealed overexpression of vesicular trafficking related genes (Rab27, Rab38, VAMPA and syntaxin 7) in the metastatic cells. Rab27 and Rab38 are small GTPases functioning at various stages of vesicular fusion and trafficking. The vesicle-associated membrane protein-associated protein A (VAMPA) and syntaxin 7 mediate the fusion of organelle membranes and vesicle attachment to

lysosomes, (Montel et al., 2005; Steeg, 2005). Overexpression of these proteins could partly explain the observed upregulation of cell surface proteins in the metastatic cells. However, when using the cell surface biotinylation method, detergents used to solubilize the membrane proteins influence the type of proteins that will be identified with the downstream applications as only the proteins soluble in the detergent will be isolated and identified. Other comparative proteomic analyses of biotinylated cell surface fractions from paired non-metastatic and metastatic cells have reported different ratios (Conn et al., 2008; Roesli et al., 2009; Luque-Garcia et al., 2010). In this thesis work, only one detergent combination (2% NP-40, 1% Triton X-100, 10% glycerol) was used and therefore many proteins insoluble in this buffer were naturally excluded from the analysis. The number of differentially expressed proteins could therefore be increased by using various different detergents, which might also affect the relative numbers of up and downregulated proteins between the cell pair.

### **5.1.3 Validation of the differential cell surface expression of selected proteins (I, unpublished data)**

In order to validate the differential expression and/or localization of the identified proteins we used Western blot and immunofluorescence analyses. In regards of the LC-MS/MS results, higher expression of PROCR (endothelial protein C receptor), CD109 and ITGA6 (integrin- $\alpha$ 6) on the surface of the metastatic cells was confirmed (Figure 3A and B in publication I). Integrin  $\beta$ 1 (ITGB1), which displayed only a 1.4-fold expression difference in the LC-MS/MS quantitation was used as a control. According to Western blot and immunofluorescence analyses it was expressed at equal levels at the cell surface of the cell pair (Figure 3A and B in publication I). In addition, the differential expression of two proteins, SSR4 and NPM, which were over- and under expressed at the metastatic cell surface in the 2D analysis, respectively, was validated in the cell surface samples. (Figure 5A, Karhemo et al, unpublished results).



**Figure 5.** Analysis of SSR and NPM expression in the non-metastatic and metastatic cells. A) Expression of SSR4 and NPM in the metastatic and non-metastatic cells was detected using Western blot analysis of cell surface extracts. Expression of NPM was also analyzed from the unbound fraction representing intracellular NPM levels. B) Immunofluorescence analysis of NPM using antibodies against NPM followed by an Alexa-594 conjugated secondary antibody. Non-metastatic and metastatic cells showed equal nuclear NPM staining. However, the metastatic cells displayed an additional cytoplasmic staining in some cells. Nuclei were visualized with DAPI (blue). Images were digitally cropped in Photoshop CS4.

Since the differential expression at the cell surface could result either from differences in the expression levels or from differential localization of proteins we analyzed the expression of CD109 and PROCR in whole cell extracts and NPM in the unbound fractions. No difference in the expression level of NPM was detected (Figure 5A, Karhemo et al., unpublished results) while both PROCR and CD109 were expressed at higher levels in the

metastatic cell line compared to the non-metastatic cells (Figure 3A in publication I). Importantly, this indicates that a change of NPM localization might play a role in metastatic colonization. Metastasis associated decrease in cell surface expression of NPM has also been reported in another cell surface proteomics study (Roesli et al., 2009). To analyze NPM cellular localization in more detail in our isogenic cell lines, we stained fixed and permeabilized cells with anti-NPM/B23 antibodies. Interestingly, in both cell lines NPM/B23 displayed strong nucleoplasmic staining and in the metastatic cell line an additional cytoplasmic localization was also observed (Figure 5B, Karhemo et al., unpublished results). As seen from Figure 5A, only a minor fraction of total cellular NPM was observed at the cell surface and is most likely below the detection limit of immunofluorescence.

#### **5.1.4 The overexpression of cd109 and ITGA6 marks cells derived from melanoma metastasis (I)**

The 3-dimensional (3D) architecture provided by the ECM at both primary and secondary sites, have a profound influence on the functional properties of tumor cells (Joyce and Pollard, 2009; Sleeman et al., 2012). Importantly, the *in vitro* observed expression differences of CD109, PTGFRN, PROCR, and ITGA6 (table 3) were also detectable *in vivo* in xenograft tumor samples derived from the metastatic and non-metastatic MDA-MB-435 cells (Figure 4 in publication I).

To investigate if the differential expression of CD109 and ITGA6 could differentiate metastatic cells from non-metastatic ones in other cell models, their expression was studied in six melanoma cell lines (WM164, WM165, WM793, WM852, G361, and Bowes) and in four invasive (MDA-MB-231, BT549, HS578T, SUM159) and four non-invasive (MCF-7, ZR75-1, BT474, and T47D) breast cancer cell lines (Neve et al., 2006). CD109 is a GPI-linked cell surface protein, which negatively modulates TGF $\beta$ 1 signaling in keratinocytes (Finnson et al., 2006; Hagiwara et al., 2010). Expression of its transcript has previously been linked to melanoma in a transgenic melanoma mouse model (Ohshima et al., 2010) and it is expressed in BLBC (Hasegawa

et al., 2008). Shedding of CD109 by mesotrypsin has been shown to promote the malignant growth of breast cancer cells (Hockla et al., 2010) but its role in metastasis is unknown.

ITGA6 dimerizes with ITGB4 and the complex functions primarily as an adhesion receptor in normal epithelia, often as a component of adhesive structures termed hemidesmosomes. The A6B4 complex has been shown to play a role in migration, invasion and survival of carcinoma cells, most likely by alterations in the localization of A6B4 that influence its signaling capacity (Lipscomb and Mercurio, 2005).

In line with the original proteomic study, CD109 was highly expressed on the surface of the cell lines derived from metastatic melanoma (WM164, WM165, and WM852) and on the surface of WM793 cells derived from an invasive advanced vertical growth phase of melanoma. Cells derived from non-metastatic primary melanomas (G361 and Bowes) expressed negligible amounts of the protein. ITGA6 was expressed in WM852 and WM793 cells while the rest of the cell lines showed very low or undetectable levels of ITGA6 (Figure 5 in publication I, control staining in Supplemental Figure S2A in publication I). In accordance with its expression in the melanoma cell lines, three out of four invasive breast cancer cell lines (BT549, HS578T, and SUM159) expressed high levels of CD109 while none of the non-invasive breast cancer cell lines expressed detectable amounts of CD109 (Supplemental Figure S1 in publication I).

To typify the fibrin-rich tumor–stroma interface of melanomas *in vivo*, the expression of CD109 and ITGA6 was further analyzed in three-dimensional (3D) matrix composed of cross-linked fibrin (Wojtukiewicz et al., 1990). Importantly, when the cells were implanted inside a 3D matrix, both ITGA6 and CD109 were highly expressed in the cells derived from melanoma metastasis (Figure 6 in publication I, control staining in Supplemental Figure S2D in publication I). WM793 cells from advanced vertical growth phase primary melanoma also expressed detectable levels of CD109 in the 3D culture conditions while ITGA6 was essentially undetectable in all the cells derived from primary melanomas (Figure 6 in publication I).

Importantly, these results demonstrate the validity of the used cell line model and the proteomics approach. The results were validated *in vivo* and in models other than originally used. Therefore, the functional role of C109 and other identified proteins in metastatic colonization and dormancy should be studied in the future.

#### **5.1.5 Interactome analysis indicates a role for a tetraspanin protein complex in metastasis (I)**

The role of only about one-third of the significantly differentially expressed proteins identified in the gel free LC-MS/MS analysis (23 proteins) is well documented in cancer progression and metastasis. To put our novel metastasis-associated proteins into biological context we analyzed how the proteins identified in the LC-MS/MS analysis (Table 3 and Table 1 in I) interacted with each other, and which signaling pathways they were involved in using the computational platform Moksiskaan (Laakso and Hautaniemi, 2010). A two-fold expression difference was used as a threshold for up or down regulated proteins. Proteins that showed less than 2-fold expression differences were defined as stable (Table 1 in I and Table 4). To simplify the pathways and interactomes the HLA-molecules were excluded from the analyses resulting in 14 differentially expressed proteins and seven stably expressed proteins. Differential expression of HLA-molecules in this cell model has been validated in other studies (Leth-Larsen et al., 2009; Rasmussen and Ditzel, 2009).

The known relationships between the candidate proteins were analyzed in the Moksiskaan software based on the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways (Figure 2 in publication I) (Kanehisa et al., 2010). The KEGG pathways supporting these relationships included metastasis-associated pathways, such as gap junctions, focal adhesions, and insulin signaling pathway (Supplemental information S2, metastasis associated proteins p. 5 in study I).

The insulin signaling pathway is represented in the analyses via two proteins upregulated at the surface of the metastatic cells; fatty acid synthase



(FASN) and receptor-type tyrosine-protein phosphatase F (PTPRF, LAR). Excess cellular lipid level has previously been associated with cancer aggressiveness (Tsubura et al., 2009). Hedegaard et al. demonstrated higher unsaturated fatty acid levels in the metastatic MDA-MB-435 cell clone relative to the non-metastatic one (Hedegaard et al., 2010). A comparative gene expression analysis revealed significantly higher expression of fatty acid desaturase 1, which catalyzes the change of saturated fatty acids to unsaturated ones, in the metastatic MDA-MB-435 cells compared to the non-metastatic ones (Montel et al., 2006). This evidence indicates a role of fatty acids in the regulation of the metastatic potential of this cell pair.

PTPRF is a protein tyrosine phosphatase (PTP) that has been reported to regulate insulin receptor phosphorylation and signaling (Goldstein et al., 1998). The extracellular region of PTPRF contains a cell adhesion molecule-like receptor region and it localizes at focal adhesions (Brady-Kalnay and Tonks, 1995), which are important for cell-extracellular matrix interactions and adhesion-mediated signal transduction (Wozniak et al., 2004). PTPs can either promote or suppress tumor progression and metastasis via either enhancing or suppressing cell surface receptor tyrosine kinase signaling (Sastry and Elferink, 2011). Upregulation of different forms of PTPs have been identified on the surface of the metastatic cells in other studies (Conn et al., 2008; Roesli et al., 2009). Interestingly, PTPRF has been shown to regulate adhesion between *Drosophila* male germ line stem cells and the niche by promoting E-cadherin-based adhesion (Srinivasan et al., 2012). It's possible role in promoting metastasis via adhesive contacts at the secondary site or pre-metastatic niche warrants further studies.

When analyzing the protein interactions in the candidate pathway we observed that three of the identified proteins, ITGA6, prostaglandin receptor negative regulator (PTGFRN) and immunoglobulin superfamily member 8 (IGSF8) interacted with two proteins (CD81 and TSPAN4) in the Moksiskaan pathway (Supplemental information S2, metastasis associated proteins p. 16 in publication I). Interestingly, CD81 and TSPAN4 are members of the tetraspanin family, a class of 33 proteins, which all have four transmembrane domains with short intracytosolic N- and C-terminal regions, and two

extracellular loops. Tetraspanins form specialized membrane microdomains on the cell surface, which control cell proliferation and migration through various adhesion and growth factor receptors. Tetraspanins interact with many different molecules, like integrins, and the composition of the complex, known as the tetraspanin web, is altered by various biological stimuli. Therefore, the different combinations of tetraspanins and associated proteins are biologically more important than individual components present at the cell surface. Recent extensive research has shown that expression of various tetraspanins and their associated partners is deregulated in human malignancies (Lazo, 2007). In regards to the PTGFRN and IGSF8, which were upregulated at the surface of the metastatic cells, it would be valuable to analyze the composition of the tetraspanin complex instead of the single proteins to better understand their role in the regulation of metastatic process.

## **5.2 Nucleophosmin (NPM/B23) has multiple divergent roles in breast cancer (II, III)**

### **5.2.1 Discovery and characterization of novel NPM splice variants B23.3 and B23.4 (III)**

As shown in Figure 5B we discovered that the metastatic as well as the non-metastatic cells both displayed strong nucleoplasmic NPM staining. In accordance, no overall expression difference of NPM was detected (Figure 5A). However, the metastatic cells lacked NPM on their surface and on the other hand contained cytoplasmic NPM (Figure 4A and B, respectively). Taken together these results indicated that a change in NPM localization might play a role in metastasis (Karhemo et al, unpublished data). NPM mutations are the most frequent genetic abnormalities found in acute myeloid leukemias and cause a cytoplasmic delocalization of the mutant NPM protein (Falini et al., 2005). To investigate the reason for the differential localization of NPM in the metastatic cells we cloned the gene from both metastatic and non-metastatic cells and analyzed its sequence.

Although no mutations in the sequences were found the sequence analysis revealed a possible explanation for the localization differences; the presence of two novel NPM splice variants in these cells. The domain structure and sequence alignment of the novel variants are shown in Figure 1 in publication III. One of the variants, hereafter referred to as B23.3 has recently been described in the human EST database (GenBank accession number: NM\_199185). The other variant, hereafter referred to as B23.4, has not been described in the human EST database.

Characterization of the cellular localization and dimerization of the novel NPM splice variants, B23.3 and B23.4 in U2OS cells, revealed that the B23.4 could be responsible for the cytoplasmic NPM observed in the metastatic cells from which it was originally identified (Figure 4G and I in publication III). According to the immunoprecipitation analysis, B23.4 did not oligomerize with the endogenous NPM (Figure 2 in III). This is not surprising taken into account that B23.4 lacked part of the oligomerization site at the N-terminus of the NPM-protein (Figure 1 in III).

B23.3 localized to the nucleoli and nucleoplasm (Figure 4D and F and 5D and F in publication III) and oligomerized with the endogenous NPM (Figure 2 in publication III). Interestingly, different NPM forms were detected in cells overexpressing the full-length NPM (B23.1) or B23.3 indicating that they formed distinct complexes in the cell (Figure 2 in publication III). The nature of these complexes is currently unknown but the observation is important because the cellular function of NPM is heavily affected by its modifications together with homo- and hetero-oligomers. Overexpression of neither of the two novel splice variants affected proliferation of U2OS cells (Figure 3A in publication III) but both affected their phenotype and post-confluent growth (Figure 3B in III). The observed phenotypic differences were also reflected in the actin filaments visualized by phalloidin staining (Figure 3C in III).

Our results obtained from the initial characterization of the novel NPM splice variants indicated that these variants are functional and therefore most likely expressed also at protein level in the cells. NPM has been shown to play diverse cellular roles depending on the cell type. Therefore, to better

understand the biological significance of these novel splice variants it will be important to study them in other cell types in addition to the U2OS cells.

Our discovery of novel NPM splice variants in addition to the previously characterized B23.2, which lacks 35 amino acids at the C-terminus (Wang et al., 1993) adds another layer of complexity to NPM's biology and highlights the need to interpret the results obtained from NPM studies in respect to not just the localization or oligomerization but in regards to the isoforms. All the isoforms seem to either lack highly specific functional domains (B23.2 and B23.3) or only contain certain domains (B23.4) indicating that the various and often seemingly antagonistic functions of NPM might regulated by its functional domains. To shed more light on the biological significance of the novel NPM splice variants their expression pattern in normal tissues under various developmental stages and in cancer tissues of multiple origin should be analyzed.

### **5.2.2 Analysis of the prognostic significance of different NPM forms in breast cancer (II, III)**

One approach to validate the relevance of observations made in experimental models is to analyze patient-derived material by using TMAs. Furthermore, such studies may identify candidate biomarkers for functional investigation. Despite the vast literature on NPM, to our knowledge, no analysis of the relevance of its expression or localization in regard to breast cancer prognosis was performed. Zhu et al. (2006) analyzed the nuclear expression of NPM in 44 breast cancer patient revealing that in 95% of cases (42/44) >50% of tumor cell nuclei were positive for NPM. However, the differences in intensity or prognostic value were not analyzed (Zhu et al., 2006).

To reveal the significance of NPM localization and/or expression level in breast cancer we investigated its expression level and localization in a large array of patient material (n =1160), the FinProg breast cancer database (Joensuu et al., 2003). To reveal the possible role of B23.3 , which lacked an important Thr199 phosphorylation site, in breast cancer we analyzed expression of the Thr199 phosphorylation (pThr199) in the same FinProg

database. To obtain a more comprehensive view a second breast cancer database, the FinHer cohort (Joensuu et al., 2009; Joensuu et al., 2006), was also analyzed for pThr199 (n=857). Both databases contain detailed clinical information and follow-up data for the patients enabling survival analysis and correlation with other known prognostic markers for NPM and its Thr199 phosphorylation. We also analyzed 14 histologically normal breast tissue samples originating from breast reduction surgery for their NPM expression for the comparison of NPM expression in breast cancer. Results from these studies are summarized in table 4. and discussed below.

**Table 4.** Different NPM forms analyzed in this study

<b>NPM form</b>	<b>Observed in</b>	<b>Prognostic factor</b>	<b>Independent prognostic factor (subtype)</b>	<b>More frequent in</b>	<b>Study</b>
<b>Cell surface</b>	NM-435	ND	ND	ND	
<b>Cyto-plasmic</b>	M-435, BCPM	No	ND	ND	I
<b>B23.3</b>	435	ND	ND	ND	III
<b>B23.4</b>	M-435	ND	ND	ND	III
<b>High NPM levels</b>	BCPM	Associate with with FP	Yes (luminal A)	No	II
<b>Granular NPM</b>	BCPM	Associate with PP	Yes (whole material, adjuvant chemotherapy)	Basal-like, unclassified	II
<b>pThr199</b>	BCPM	Associate with PP	No	Basal-like	III
<b>NM-435 = non-metastatic MDA-MB-435 subclone,  M-435 = metastatic MDA-MB-435 subclone,  BCPM = breast cancer patient material,  ND = not determined, FP = favourable protgnosis, PP = poor prognosis</b>					

### 5.2.3 NPM expression levels affected its localization (II)

In our analysis, NPM was expressed at high levels in the luminal epithelial cells of all histologically normal breast tissue samples (Figure 1A, B and E in publication II). Similar intense, nuclear staining of the epithelial cells of the milk ducts has been detected in another study (Rower et al., 2011) indicating that under normal conditions, the luminal epithelial cells have high levels of total NPM. Interestingly, in our histologically normal breast tissue samples the younger women displayed uniform nuclear NPM localization (Figure 1A in publication II) while the older ones showed only nucleolar localization (Figure 1B in publication II). By using qPCR analysis and ectopic overexpression of NPM we showed that NPM expression level affected its localization (nucleolar or nucleoplasmic, Figure 1G-I in publication II). We confirmed the previously published results (Cicatiello et al., 2004; Skaar et al., 1998) that NPM expression is up-regulated by estrogen treatment in breast cancer cell lines (Figure 2A and B in publication) and showed that this upregulation leads to increase in the nucleoplasmic NPM (Figure 2C-F in publication II). Since estrogen levels commonly decrease at menopause in older women our results suggest that NPM expression level affected its localization in the. In accordance with our findings, UV treatment (stress) induces up-regulation of NPM expression and changes its localization from nucleolar to even nuclear distribution (Wu and Yung, 2002). If and how the effects of the stress-induced nucleoplasmic localization of NPM (Chan et al., 1985; Yang et al., 2002; Kurki et al., 2004) differ from the nucleoplasmic localization in the breast epithelial cells is unknown.

In the breast cancer samples analyzed, NPM was detected at various localizations: solely in the nucleolus (14%, Figure 3A, F in II), evenly distributed throughout the nucleus (55%, Figure 3 B, G in II), and only in the nucleoplasm (31%, Figure 3 C, H in II). We also observed a rare, cytoplasmic staining for NPM (7%, Figure 3 D, I in II). However, no correlation between the patient prognosis and these NPM localizations was observed in the study. In contrast to histologically normal breast tissue, no correlation between the NPM localization and age was detected in the tumor specimens ( $P = 0.67$ )

and no mutations explaining the differences were found (Supplemental table S1 in publication II).

#### **5.2.4 Tumor suppressive role for NPM in breast cancer (II)**

Importantly, NPM expression was reduced in breast cancer as compared to histologically normal breast tissue and patients with reduced levels of NPM showed increased risk of developing a metastatic disease as judged by distant disease free survival (DDFS, time from diagnosis to the occurrence of either metastases outside the region in which the tumor arose or death from breast cancer) (Figure 4A and supplemental Table S3 in publication II). In a subgroup analysis NPM levels showed significant prognostic value in ER-positive, p53<sup>-/low</sup> and HER-negative breast cancers. More importantly, NPM expression levels served as an independent prognostic factor for metastatic disease and/or breast cancer caused death in the luminal A subgroup of breast cancer (Figure 4B in II and Table 2 in II). The luminal A tumors in our analysis were classified as ER-and/or PR-positive and HER-negative, which explains the observed prognostic value in these subgroups. The prognostic value in p53<sup>-/low</sup>-groups, which most likely have a wild type p53, can be explained by the knowledge that only about 12-15% of luminal A tumors harbor p53 mutations (Carey et al., 2006; Cancer Genome Atlas Network, 2012). Luminal B tumors differ from the luminal A ones in respect to HER2 expression. NPM expression showed no prognostic value in this subtype or in HER2-positive tumors indicating that HER2 overexpression overrode the prognostic significance of NPM.

Our results revealed that overexpression of NPM in the invasive MDA-MB-231 cells abrogated their growth in soft agar (Figure 5 in II) suggesting a tumor suppressive role for NPM in breast cancer. Contradictory to our results, high NPM expression has been associated with poor prognosis or recurrence in other tumor types (Kikuta et al., 2009; Tsui et al., 2008; Liu et al., 2012; Coutinho-Camillo et al., 2010). However, in the cases where NPM expression in the corresponding normal tissue was analyzed, it was markedly lower than in the cancer tissue highlighting the context dependent role for

NPM in neoplasia. Too much or too little NPM could disrupt the cellular homeostasis leading to deregulated growth.

#### **5.2.5 Granular staining pattern as indicator of metastatic disease (II)**

Interestingly, 14% of the breast cancer samples showed a very distinct, granular staining for NPM (Figure 3E and J in publication II). A similar staining pattern has previously been identified in colorectal carcinoma (Nozawa et al., 1996) but its significance has not been analyzed. In this thesis work the presence of the granular staining pattern associated with aggressive breast cancer characters and basal subtype (Supplemental Table S2 in publication II). Moreover, patients that showed granular NPM were more prone to develop metastatic disease or die of breast cancer than patients with non-granular NPM (Figure 4C in publication II) in several subgroups (Supplemental Table S4 in II) including luminal A (Figure 4D in II). More importantly, in a Cox proportional hazards model the granular staining pattern proved as an independent prognostic factor of distant, metastatic disease or breast cancer death in the whole series (Table 3 in II,  $P = 0.04$ ). Patients displaying granular staining pattern of NPM appeared less likely to benefit from chemotherapy (Figure 4, E and F in II and Table 4 in publication II). The cause of the granular NPM is currently unknown but the mechanisms behind it would be interesting to resolve since they might shed light on the contradiction between the oncogenic and tumor suppressive functions of NPM.

#### **5.2.6 Thr199 phosphorylation of NPM as an indicator of recurrence (III)**

Interestingly, Thr199 phosphorylation (pThr199) of NPM associated with development of distant, metastatic disease and death from breast cancer (DDFS,  $P=0.016$ , Figure 7A in III) in the NPM positive (moderate or high expression) breast cancers in the FinProg dataset. This result suggests that Thr199 phosphorylation of NPM overrides the tumor suppressive effect



observed for the high total NPM levels and that NPMpThr199 represents an oncogenic form of the NPM protein. In support of this, pThr199 significantly associated with poor overall survival ( $P=0,013$ , Figure 7B in publication II) and distant disease recurrence, in other words metastatic disease ( $P=0,008$ , Figure 7C in publication II) in the FinHer material that contains patients with more aggressive disease. In this respect, the NPMpThr199 resembled the granular staining pattern. Both NPM forms were also more frequently observed in basal type breast cancers. These two forms of NPM showed inverse correlation in the FinProg breast cancer patients (Table 1 in III) indicating a connection between them. The nature of this connection, however, is totally unclear. Since nothing is known about the molecular basis of granular NPM, it is impossible to speculate the underlining mechanisms behind the similar behavior and connection between the granular NPM and Thr199 phosphorylation of NPM.

#### **5.2.7 Thr199 phosphorylation of NPM associated with aggressive breast cancers and CIP2A expression (III)**

Cancerous inhibitor of protein phosphatase 2 (CIP2A) has recently been identified as an oncogene (Junttila et al., 2007). Intriguingly, we demonstrated that NPMp Thr199 associated with CIP2A expression in both breast cancer cohorts analyzed (Table 1 in III and Figure 6A in publication III). Moreover, a significant reduction in pThr199 levels was observed at day 4 and 5 after siRNA mediated silencing of CIP2A (Figure 6B in III) confirming the association observed in the patient analysis. CIP2A also seemed to associate with total NPM levels in the patient material. In accordance, a slight increase in total NPM levels was evident at day 3 after CIP2A silencing, but this was not detected at later time points (Figure 6B in III).

Thr199 of NPM is dephosphorylated by a protease distinct from PP2A, PP1beta (Lin et al., 2010). In addition, CIP2A silencing increased the pThr199 at day 3 after which a clear reduction was observed (Figure 6B in

III). This together with the different cellular localizations of CIP2A and NPM indicate that Thr199 is not a direct target of PP2A.

The expression of CIP2A (Côme et al., 2009; Niemela et al., 2012) and the NPMpThr199 both associated with aggressive characteristics of breast cancer (p53 expression indicative of mutant p53 and basal-type, Table 1 in publication III). These two proteins participate in the regulation of the same pathways like c-myc (Niemela et al., 2012; Bockelman et al., 2012; Guo et al., 2000; Neiman et al., 2001; Kim et al., 2000; Zeller et al., 2001; Li and Hann, 2013; Li et al., 2008) and E2F1 (Lin et al., 2010; Lin et al., 2006; Takemura et al., 2002; Takemura et al., 1999; Liu et al., 2009; Laine et al., 2013). In addition, in our analysis, pThr199 significantly associated with EGFR expression (Table 1 in publication III) and the CIP2A has shown to be regulated by the EGFR-MEK1/2 ETS signaling pathway (Khanna et al., 2011). The detailed molecular mechanism behind the connection between NPMpThr199 and CIP2A warrants further experimental analyses to resolve if CIP2A is responsible for the association between NPMpThr199 and development of metastatic disease in breast cancer.

#### **5.2.8 NPM modifications and metastasis?**

In this thesis, by using two distinct large breast cancer TMA datasets, we have revealed that, in respect of development of metastatic disease and breast cancer death, different NPM forms have contradictory roles in breast cancer (summarized in Table 4). To our knowledge, this is the first analysis comparing the significance of different NPM forms (total levels, granular NPM and Thr199 phosphorylation) in any cancer material. Interestingly, high total NPM levels associated with favourable prognosis and reduced risk of metastatic disease and/or death from breast cancer only in luminal A breast tumors while its expression did not significantly vary among the different subtypes (Supplemental Table S2 in publication II). On the contrary, granular staining and Thr199 phosphorylation were more frequent in basal-like breast cancers and served as indicators of metastatic disease in whole patient material. As described in the Review of the literature, NPM

participates in the regulation of p53 and RB-pathways. Interestingly, vast majority of luminal A tumors harbor wt p53 while this protein is mutated or inactivated in most BLBCs (Carey et al., 2006; Cancer Genome Atlas Network, 2012; Dumay et al., 2013). In addition, in most BLBCs tumor suppressor RB is affected (Gauthier et al., 2007; Herschkowitz et al., 2008; Subhawong et al., 2009). If these molecular differences affect the role of NPM and its different forms in breast cancer should be analyzed in the future.

Previously, in a comparative proteomic analysis the phosphorylation of NPM serine 125 has been shown to be more abundant in cancer samples than in controls (Rower et al., 2011). In addition, a 2D gel analysis identified an unknown NPM modification to be more abundant in metastatic breast tumors than in their non-metastatic counterparts (Vydra et al., 2008). Moreover, this thesis works revealed a novel NPM splice variant from metastatic cells and a connection between NPMpThr199 and a known oncogene and modulator of post-translational modifications, CIP2A. Taken together these findings propose that postranslational modifications and domain structure changes affecting these modifications in NPM might be general mechanisms to control NPM's function in breast cancer. To make the picture more complicated, NPM's oncogenic potential is also modified by plakoglobin (PG,  $\gamma$ -catenin), a homolog of  $\beta$ -catenin with dual adhesive and signaling functions (Lam et al., 2012) and HLJ1, a tumor suppressor and a member of the heat shock protein 40 chaperone family (Chang et al., 2010). In addition, NPM has been shown to suppress of CXCR4-mediated G protein activation and chemotaxis via directly interacting with CXCR4. CXCR4 is the primary receptor for CXCL12, which is secreted by metastatic sites and promotes homing of CXCR4 expressing tumor cells to these sites (Zhang et al., 2007).

## 6 CONCLUSIONS AND FUTURE PROSPECTS

One aim of this thesis work was to identify novel cell surface proteins associated with metastasis colonization and dormancy. Importantly, this study resulted in the discovery of a several novel metastasis associated proteins and opened up interesting connections for further studies. Out of the identified proteins, CD109 proved as a promising novel candidate for future analyses. In addition to the conventional cell surface proteins the proteomic comparison revealed novel cell surface proteins. Intracellular proteins are increasingly observed in extracellular space highlighting the need to reconsider the classical protein localizations. Importantly, in respect to drug target discovery a protein that is a therapeutic target in one cellular location may have anti-target activity elsewhere.

The present study demonstrated for the first time analysis of different NPM forms in the same patient material. Strikingly, different NPM forms (total levels, granular staining and pThr199) behaved in opposite ways indicating the need to critically evaluate results concerning NPM's function and prognostic relevance. NPM cannot be considered as a single entity but in the future more effort should be put to analyses to study if similar relationships are present in other tumor types. In respect to NPM as a drug target one should carefully consider all the possible effects the drug might have on NPM's function.

The pThr199 associated with CIP2A expression. The positive prognostic role of NPM was evident in luminal A tumors, which rarely harbor p53 mutations. The poor prognostic function was evident in basal type breast cancer commonly harboring a mutant p53. The connection of NPM to p53 and CIP2A should be analyzed at functional level.

Finally, this study revealed the presence of two novels splice variants of NPM, B23.3 and B23.4, in cancer cells. In the future their molecular function, expression pattern at the tissue level and possible role in regulation of metastatic dissemination of cancer should be studied.

## 7 ACKNOWLEDGEMENTS

*This work was carried out at the University of Helsinki, Faculty of Medicine, Research Programs Unit, Molecular Cancer Biology Research Program and its successor Translational Cancer Biology Research Program and at the institute of Biomedicine. I wish to thank Professors Olli Jänne, Sampsa Hautaniemi, Marikki Laiho, Jorma Keski-Oja, Kari Alitalo and Esa Korpi for providing the great research facilities in the Biomedicum Helsinki.*

*I am greatly thankful to my supervisor Docent Pirjo Laakkonen for not only giving me the opportunity to work in your laboratory but especially for the possibility to work with such an interesting and challenging project. You helped me to develop in my scientific skills and gave me the opportunity to test my own ideas. I have enjoyed our fruitful discussions and debates. You certainly have made me feel like an appreciated member of your team.*

*Docent Markku Varjosalo and Professor Jorma Isola are greatly acknowledged for their thoughtful review and constructive comments on how to improvements this thesis. I also wish to thank my thesis committee members Päivi Ojala and Tuula Nyman for their valuable advises and encouragement during our annual meetings.*

*I express my sincere gratitude for Professor Jonathan Sleeman for accepting the assignment as the opponent of my thesis.*

*The past and present members of the Laakkonen lab, Minna, Juulia, Päivi, Antti, Johanna, Selina, Anastasiya, Pauliina, Maija- Thank you all for helping me so much with the experiments. Special thanks to Juulia with whom we set up many practices in the Laakkonen lab as the first two graduate students and shared many immemorial moments in and out of the lab. Maija, I appreciate your support, our discussions and especially the babysitting in the lab during late manuscript submission hours. Antti, your help with the nucleophosmin work was invaluable.*

*I thank all the co-authors of the original publications. Harri Sihto is thanked for nice and fruitful discussions, advises and analyses, Heikki Joensuu, Petri Bono and Päivi Heikkilä for the clinical material and Johan*

*and Mikael Lundin for statistical analyses. Suvi, Leena, and Ilja, thank you for your help with the proteomics paper. Kaisa and Olga were indispensable help in finalizing the proteomics study. Marko and Sampsa are acknowledged for the interactome and pathway analyses, Anni Laine and Jukka Westermarck for sharing their tools and knowledge concerning CIP2A and Marikki Laiho and Henna Moore for fruitful discussions, advises and collaboration concerning the NPM splice variants. Kimmo Porkka and Marc Baumann are thanked for their valuable support in the early phases of this thesis work. Tapio Tainola is acknowledged for his excellent management skills, guidance and advises and the Biomedicum Imaging unit for help with the microscopes.*

*I thank previous members of the Alitalo lab (Jarkko, Hanski, Sanna, Hanna, Maria, Katja and others) for creating a wonderful working environment and for activities outside the lab. Karita and Henna from Laiho lab and Anita from Saharinen Lab, thank you for encouraging me at the frustrating moments when nothing seemed to go right. I also want to thank all my friends outside the lab, especially Natasha and Katja.*

*I thank my parents Pirjo and Hannu for being there when ever needed and my sisters Anna and Suvi for their support along the way. I thank my husband Kimmo for his love and for putting things into perspective and my precious daughter, Amanda, for teaching me the skill to leave the science and enjoy other important aspects in life.*

*I wish to thank the Helsinki Graduate Program in Biotechnology and Molecular Biology for their financial support as well as for organizing interesting courses. This study has also been supported by grants from the Academy of Finland, the European Union, veritautien tutkimussäätiö, The University of Helsinki Fund, the K. Albin Johansson Foundation, Maud Kuistila Memorial Foundation, Magnush Ehnrooth foundation, Ida Montin foundation, Oskar Öfflund foundation and Otto A Malmi foundation.*

Helsinki, May 2013

## 8 REFERENCES

- Abdel-Ghany M, Cheng HC, Elble RC, Pauli BU. The breast cancer beta 4 integrin and endothelial human CLCA2 mediate lung metastasis. *J Biol Chem*. 2001 Jul 6;276(27):25438-46.
- Adon AM, Zeng X, Harrison MK, Sannem S, Kiyokawa H, Kaldis P, et al. Cdk2 and Cdk4 regulate the centrosome cycle and are critical mediators of centrosome amplification in p53-null cells. *Mol Cell Biol*. 2010 Feb;30(3):694-710.
- Aguirre Ghiso JA, Kovalski K, Ossowski L. Tumor dormancy induced by downregulation of urokinase receptor in human carcinoma involves integrin and MAPK signaling. *J Cell Biol*. 1999 Oct 4;147(1):89-104.
- Aguirre-Ghiso JA, Liu D, Mignatti A, Kovalski K, Ossowski L. Urokinase receptor and fibronectin regulate the ERK(MAPK) to p38(MAPK) activity ratios that determine carcinoma cell proliferation or dormancy in vivo. *Mol Biol Cell*. 2001 Apr;12(4):863-79.
- Albiero E, Madeo D, Bolli N, Giaretta I, Bona ED, Martelli MF, et al. Identification and functional characterization of a cytoplasmic nucleophosmin leukaemic mutant generated by a novel exon-11 NPM1 mutation. *Leukemia*. 2007 May;21(5):1099-103.
- Allan AL, Vantyghem SA, Tuck AB, Chambers AF. Tumor dormancy and cancer stem cells: Implications for the biology and treatment of breast cancer metastasis. *Breast Dis*. 2006 - 2007;26:87-98.
- Anderson WF, Reeves JE, Elias A, Berkel H. Outcome of patients with metastatic breast carcinoma treated at a private medical oncology clinic. *Cancer*. 2000 Jan 1;88(1):95-107.
- Arvold ND, Taghian AG, Niemierko A, Abi Raad RF, Sreedhara M, Nguyen PL, et al. Age, breast cancer subtype approximation, and local recurrence after breast-conserving therapy. *J Clin Oncol*. 2011 Oct 10;29(29):3885-91.
- Ashcroft M, Vousden KH. Regulation of p53 stability. *Oncogene*. 1999 Dec 13;18(53):7637-43.
- Atchley DP, Albarracin CT, Lopez A, Valero V, Amos CI, Gonzalez-Angulo AM, et al. Clinical and pathologic characteristics of patients with BRCA-positive and BRCA-negative breast cancer. *J Clin Oncol*. 2008 Sep 10;26(26):4282-8.
- Auerbach R, Lu WC, Pardon E, Gumkowski F, Kaminska G, Kaminski M. Specificity of adhesion between murine tumor cells and capillary endothelium: An in vitro correlate of preferential metastasis in vivo. *Cancer Res*. 1987 Mar 15;47(6):1492-6.
- Avninder S, Ylaya K, Hewitt SM. Tissue microarray: A simple technology that has revolutionized research in pathology. *J Postgrad Med*. 2008 Apr-Jun;54(2):158-62.
- Barkan D, El Touny LH, Michalowski AM, Smith JA, Chu I, Davis AS, et al. Metastatic growth from dormant cells induced by a col-I-enriched fibrotic environment. *Cancer Res*. 2010 Jul 15;70(14):5706-16.
- Barkan D, Green JE, Chambers AF. Extracellular matrix: A gatekeeper in the transition from dormancy to metastatic growth. *Eur J Cancer*. 2010 May;46(7):1181-8.
- Beckmann R, Buchner K, Jungblut PR, Eckerskorn C, Weise C, Hilbert R, et al. Nuclear substrates of protein kinase C. *Eur J Biochem*. 1992 Nov 15;210(1):45-51.
- Bendas G, Borsig L. Cancer cell adhesion and metastasis: Selectins, integrins, and the inhibitory potential of heparins. *Int J Cell Biol*. 2012;2012:676731.

## REFERENCES

- Bernards R, Weinberg RA. A progression puzzle. *Nature*. 2002 Aug 22;418(6900):823.
- Bertwistle D, Sugimoto M, Sherr CJ. Physical and functional interactions of the arf tumor suppressor protein with nucleophosmin/B23. *Mol Cell Biol*. 2004 Feb;24(3):985-96.
- Berx G, Raspe E, Christofori G, Thiery JP, Sleeman JP. Pre-EMTing metastasis? recapitulation of morphogenetic processes in cancer. *Clin Exp Metastasis*. 2007;24(8):587-97.
- Bhat UG, Jagadeeswaran R, Halasi M, Gartel AL. Nucleophosmin interacts with FOXM1 and modulates the level and localization of FOXM1 in human cancer cells. *J Biol Chem*. 2011 Dec 2;286(48):41425-33.
- Blows FM, Driver KE, Schmidt MK, Broeks A, van Leeuwen FE, Wesseling J, et al. Subtyping of breast cancer by immunohistochemistry to investigate a relationship between subtype and short and long term survival: A collaborative analysis of data for 10,159 cases from 12 studies. *PLoS Med*. 2010 May 25;7(5):e1000279.
- Bockelman C, Koskensalo S, Hagstrom J, Lundin M, Ristimaki A, Haglund C. CIP2A overexpression is associated with c-myc expression in colorectal cancer. *Cancer Biol Ther*. 2012 Mar;13(5):289-95.
- Bocker T, Bittinger A, Wieland W, Buettner R, Fauser G, Hofstaedter F, et al. In vitro and ex vivo expression of nucleolar proteins B23 and p120 in benign and malignant epithelial lesions of the prostate. *Mod Pathol*. 1995 Apr;8(3):226-31.
- Bolli N, Nicoletti I, De Marco MF, Bigerna B, Pucciarini A, Mannucci R, et al. Born to be exported: COOH-terminal nuclear export signals of different strength ensure cytoplasmic accumulation of nucleophosmin leukemic mutants. *Cancer Res*. 2007 Jul 1;67(13):6230-7.
- Borer RA, Lehner CF, Eppenberger HM, Nigg EA. Major nucleolar proteins shuttle between nucleus and cytoplasm. *Cell*. 1989 Feb 10;56(3):379-90.
- Brady SN, Maggi LB, Jr, Winkeler CL, Toso EA, Gwinn AS, Pelletier CL, et al. Nucleophosmin protein expression level, but not threonine 198 phosphorylation, is essential in growth and proliferation. *Oncogene*. 2009 Sep 10;28(36):3209-20.
- Brady SN, Yu Y, Maggi LB, Jr, Weber JD. ARF impedes NPM/B23 shuttling in an Mdm2-sensitive tumor suppressor pathway. *Mol Cell Biol*. 2004 Nov;24(21):9327-38.
- Brady-Kalnay SM, Tonks NK. Protein tyrosine phosphatases as adhesion receptors. *Curr Opin Cell Biol*. 1995 Oct;7(5):650-7.
- Braun S, Vogl FD, Naume B, Janni W, Osborne MP, Coombes RC, et al. A pooled analysis of bone marrow micrometastasis in breast cancer. *N Engl J Med*. 2005 Aug 25;353(8):793-802.
- Brinkley BR, Beall PT, Wible LJ, Mace ML, Turner DS, Cailleau RM. Variations in cell form and cytoskeleton in human breast carcinoma cells in vitro. *Cancer Res*. 1980 Sep;40(9):3118-29.
- Brown DM, Ruoslahti E. Metadherin, a cell surface protein in breast tumors that mediates lung metastasis. *Cancer Cell*. 2004 Apr;5(4):365-74.
- Burgess RJ, Zhang Z. Histone chaperones in nucleosome assembly and human disease. *Nat Struct Mol Biol*. 2013 Jan;20(1):14-22.
- Burkhardt D, L., Sage J. Cellular mechanisms of tumour suppression by the retinoblastoma gene. *Nature reviews Cancer*. 2008;8(9):671-82.
- Butler TP, Gullino PM. Quantitation of cell shedding into efferent blood of mammary adenocarcinoma. *Cancer Res*. 1975 Mar;35(3):512-6.
- Cailleau R, Olive M, Cruciger QV. Long-term human breast carcinoma cell lines of metastatic origin: Preliminary characterization. *In Vitro*. 1978 Nov;14(11):911-5.



Calli AO, Sari A, Evcim G, Altinboga A, Sadullahoglu C, Ermete M. The enigmatic role of nucleophosmin in malignant melanoma: Does it have an effect? *Indian J Pathol Microbiol.* 2011 Jul-Sep;54(3):482-6.

Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature.* 2012 Oct 4;490(7418):61-70.

Cancer statistics at finnish cancer registry [Internet].: Finnish Cancer Registry. Available from: [www.cancerregistry.fi](http://www.cancerregistry.fi).

Carey L, Winer E, Viale G, Cameron D, Gianni L. Triple-negative breast cancer: Disease entity or title of convenience? *Nat Rev Clin Oncol.* 2010 Dec;7(12):683-92.

Carey LA, Perou CM, Livasy CA, Dressler LG, Cowan D, Conway K, et al. Race, breast cancer subtypes, and survival in the carolina breast cancer study. *JAMA.* 2006 Jun 7;295(21):2492-502.

Castano Z, Tracy K, McAllister SS. The tumor macroenvironment and systemic regulation of breast cancer progression. *Int J Dev Biol.* 2011;55(7-9):889-97.

Cha H, Hancock C, Dangi S, Maignel D, Carrier F, Shapiro P. Phosphorylation regulates nucleophosmin targeting to the centrosome during mitosis as detected by cross-reactive phosphorylation-specific MKK1/MKK2 antibodies. *Biochem J.* 2004 Mar 15;378(Pt 3):857-65.

Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis. *Science.* 2011 Mar 25;331(6024):1559-64.

Chambers AF. MDA-MB-435 and M14 cell lines: Identical but not M14 melanoma? *Cancer Res.* 2009 Jul 1;69(13):5292-3.

Chambers AF, Groom AC, MacDonald IC. Dissemination and growth of cancer cells in metastatic sites. *Nat Rev Cancer.* 2002 Aug;2(8):563-72.

Chan PK, Chan FY, Morris SW, Xie Z. Isolation and characterization of the human nucleophosmin/B23 (NPM) gene: Identification of the YY1 binding site at the 5' enhancer region. *Nucleic Acids Res.* 1997 Mar 15;25(6):1225-32.

Chan PK, Aldrich M, Busch H. Alterations in immunolocalization of the phosphoprotein B23 in HeLa cells during serum starvation. *Exp Cell Res.* 1985 Nov;161(1):101-10.

Chang JH, Lin JY, Wu MH, Yung BY. Evidence for the ability of nucleophosmin/B23 to bind ATP. *Biochem J.* 1998 Feb 1;329 ( Pt 3)(Pt 3):539-44.

Chang TP, Yu SL, Lin SY, Hsiao YJ, Chang GC, Yang PC, et al. Tumor suppressor HLJ1 binds and functionally alters nucleophosmin via activating enhancer binding protein 2alpha complex formation. *Cancer Res.* 2010 Feb 15;70(4):1656-67.

Chavez KJ, Garimella SV, Lipkowitz S. Triple negative breast cancer cell lines: One tool in the search for better treatment of triple negative breast cancer. *Breast Dis.* 2010;32(1-2):35-48.

Cheang MC, Chia SK, Voduc D, Gao D, Leung S, Snider J, et al. Ki67 index, HER2 status, and prognosis of patients with luminal B breast cancer. *J Natl Cancer Inst.* 2009 May 20;101(10):736-50.

Cheng HC, Abdel-Ghany M, Elble RC, Pauli BU. Lung endothelial dipeptidyl peptidase IV promotes adhesion and metastasis of rat breast cancer cells via tumor cell surface-associated fibronectin. *J Biol Chem.* 1998 Sep 11;273(37):24207-15.

Choi JW, Lee SB, Ahn JY, Lee KH. Disruption of ATP binding destabilizes NPM/B23 and inhibits anti-apoptotic function. *BMB Rep.* 2008 Dec 31;41(12):840-5.

Choi JW, Lee SB, Kim CK, Lee KH, Cho SW, Ahn JY. Lysine 263 residue of NPM/B23 is essential for regulating ATP binding and B23 stability. *FEBS Lett.* 2008 Apr 2;582(7):1073-80.

## REFERENCES

- Cicatiello L, Scafoglio C, Altucci L, Cancemi M, Natoli G, Facchiano A, et al. A genomic view of estrogen actions in human breast cancer cells by expression profiling of the hormone-responsive transcriptome. *J Mol Endocrinol*. 2004 Jun;32(3):719-75.
- Cobrinik D. Pocket proteins and cell cycle control.(cyclins and CDKs in development and cancer)(review). *Oncogene*, April 18, 2005, Vol 24(17), p 2796. 2005;24(17):2796.
- Collins VP, Loeffler RK, Tivey H. Observations on growth rates of human tumors. *Am J Roentgenol Radium Ther Nucl Med*. 1956 Nov;76(5):988-1000.
- Colombo E, Bonetti P, Lazzerini Denchi E, Martinelli P, Zamponi R, Marine JC, et al. Nucleophosmin is required for DNA integrity and p19Arf protein stability. *Mol Cell Biol*. 2005 Oct;25(20):8874-86.
- Colombo E, Marine JC, Danovi D, Falini B, Pelicci PG. Nucleophosmin regulates the stability and transcriptional activity of p53. *Nat Cell Biol*. 2002 Jul;4(7):529-33.
- Côme C, Laine A, Chanrion M, Edgren H, Mattila E, Liu X, et al. CIP2A is associated with human breast cancer aggressivity. *Clinical Cancer Research*. 2009;15(16):5092.
- Comen E, Norton L, Massague J. Clinical implications of cancer self-seeding. *Nat Rev Clin Oncol*. 2011 Jun;8(6):369-77.
- Comen EA, Norton L, Massague J. Breast cancer tumor size, nodal status, and prognosis: Biology trumps anatomy. *J Clin Oncol*. 2011 Jul 1;29(19):2610-2.
- Conn EM, Madsen MA, Cravatt BF, Ruf W, Deryugina EI, Quigley JP. Cell surface proteomics identifies molecules functionally linked to tumor cell intravasation. *J Biol Chem*. 2008 Sep 26;283(39):26518-27.
- Conn EM, Madsen MA, Cravatt BF, Ruf W, Deryugina EI, Quigley JP. Cell surface proteomics identifies molecules functionally linked to tumor cell intravasation. *J Biol Chem*. 2008 Sep 26;283(39):26518-27.
- Cordwell SJ, Thingholm TE. Technologies for plasma membrane proteomics. *Proteomics*. 2010 Feb;10(4):611-27.
- Corson TW. One hit, two hits, three hits, more? genomic changes in the development of retinoblastoma. *GENES CHROMOSOMES & CANCER*. 2007;46(7):617-34.
- Coutinho-Camillo CM, Lourenco SV, Nishimoto IN, Kowalski LP, Soares FA. Nucleophosmin, p53, and ki-67 expression patterns on an oral squamous cell carcinoma tissue microarray. *Hum Pathol*. 2010 Aug;41(8):1079-86.
- Dawood S, Hu R, Homes MD, Collins LC, Schnitt SJ, Connolly J, et al. Defining breast cancer prognosis based on molecular phenotypes: Results from a large cohort study. *Breast Cancer Res Treat*. 2011 Feb;126(1):185-92.
- den Besten W, Kuo ML, Williams RT, Sherr CJ. Myeloid leukemia-associated nucleophosmin mutants perturb p53-dependent and independent activities of the arf tumor suppressor protein. *Cell Cycle*. 2005 Nov;4(11):1593-8.
- Dhar SK, St Clair DK. Nucleophosmin blocks mitochondrial localization of p53 and apoptosis. *J Biol Chem*. 2009 Jun 12;284(24):16409-18.
- Dimaras H. Loss of RB1 induces non-proliferative retinoma: Increasing genomic instability correlates with progression to retinoblastoma. *Hum Mol Genet*. 2008;17(10):1363-72.
- Domon B, Aebersold R. Options and considerations when selecting a quantitative proteomics strategy. *Nat Biotechnol*. 2010 Jul;28(7):710-21.

- Dontu G, Al-Hajj M, Abdallah WM, Clarke MF, Wicha MS. Stem cells in normal breast development and breast cancer. *Cell Prolif*. 2003 Oct;36 Suppl 1:59-72.
- Du W, Zhou Y, Pike S, Pang Q. NPM phosphorylation stimulates Cdk1, overrides G2/M checkpoint and increases leukemic blasts in mice. *Carcinogenesis*. 2010 Feb;31(2):302-10.
- Dumay A, Feugeas JP, Wittmer E, Lehmann-Che J, Bertheau P, Espie M, et al. Distinct tumor protein p53 mutants in breast cancer subgroups. *Int J Cancer*. 2013 Mar 1;132(5):1227-31.
- Dumbar TS, Gentry GA, Olson MOJ. Interaction of nucleolar phosphoprotein B23 with nucleic acids. *Biochemistry (N Y)*. 1989;28(24):9495.
- Duncan MW, Aebersold R, Caprioli RM. The pros and cons of peptide-centric proteomics. *Nat Biotechnol*. 2010 Jul;28(7):659-64.
- Eischen CM, Weber JD, Roussel MF, Sherr CJ, Cleveland JL. Disruption of the ARF-Mdm2-p53 tumor suppressor pathway in myc-induced lymphomagenesis. *Genes Dev*. 1999 Oct 15;13(20):2658-69.
- Elia G. Biotinylation reagents for the study of cell surface proteins. *Proteomics*. 2008 Oct;8(19):4012-24.
- Ellison G, Klinowska T, Westwood RF, Docter E, French T, Fox JC. Further evidence to support the melanocytic origin of MDA-MB-435. *Mol Pathol*. 2002 Oct;55(5):294-9.
- Elschenbroich S, Kim Y, Medin JA, Kislinger T. Isolation of cell surface proteins for mass spectrometry-based proteomics. *Expert Rev Proteomics*. 2010 Feb;7(1):141-54.
- Elstrodt F, Hollestelle A, Nagel JH, Gorin M, Wasielewski M, van den Ouweland A, et al. BRCA1 mutation analysis of 41 human breast cancer cell lines reveals three new deleterious mutants. *Cancer Res*. 2006 Jan 1;66(1):41-5.
- Ewing J. Neoplastic diseases: A treatise on tumors. 3rd ed ed. Philadelphia (PA): WB Saunders; 1928.
- Falini B, Bolli N, Liso A, Martelli MP, Mannucci R, Pileri S, et al. Altered nucleophosmin transport in acute myeloid leukaemia with mutated NPM1: Molecular basis and clinical implications. *Leukemia*. 2009 Oct;23(10):1731-43.
- Falini B, Nicoletti I, Martelli MF, Mecucci C. Acute myeloid leukemia carrying cytoplasmic/mutated nucleophosmin (NPMc+ AML): Biologic and clinical features. *Blood*. 2007 Feb 1;109(3):874-85.
- Falini B, Mecucci C, Tiacci E, Alcalay M, Rosati R, Pasqualucci L, et al. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med*. 2005 Jan 20;352(3):254-66.
- Federici L, Arcovito A, Scaglione GL, Scaloni F, Lo Sterzo C, Di Matteo A, et al. Nucleophosmin C-terminal leukemia-associated domain interacts with G-rich quadruplex forming DNA. *J Biol Chem*. 2010 Nov 26;285(48):37138-49.
- Ferlay J, Shin H, Bray F, Forman D, Mathers C, Parkin D. GLOBOCAN 2008 v1.2, cancer incidence and mortality worldwide: IARC CAnceRBase no. 10 [<http://Globocan.iarc.fr>]. . 2011.
- Feuerstein N, Chan PK, Mond JJ. Identification of numatrin, the nuclear matrix protein associated with induction of mitogenesis, as the nucleolar protein B23. implication for the role of the nucleolus in early transduction of mitogenic signals. *J Biol Chem*. 1988 Aug 5;263(22):10608-12.
- Feuerstein N, Spiegel S, Mond JJ. The nuclear matrix protein, numatrin (B23), is associated with growth factor-induced mitogenesis in swiss 3T3 fibroblasts and with T lymphocyte proliferation

## REFERENCES

- stimulated by lectins and anti-T cell antigen receptor antibody. *J Cell Biol.* 1988 Nov;107(5):1629-42.
- Feuerstein N, Mond JJ. "Numatrin," a nuclear matrix protein associated with induction of proliferation in B lymphocytes. *J Biol Chem.* 1987 Aug 15;262(23):11389-97.
- Feuerstein N, Mond JJ. Identification of a prominent nuclear protein associated with proliferation of normal and malignant B cells. *J Immunol.* 1987 Sep 15;139(6):1818-22.
- Fidler IJ. The pathogenesis of cancer metastasis: The 'seed and soil' hypothesis revisited. *Nat Rev Cancer.* 2003 Jun;3(6):453-8.
- Fidler IJ, Kripke ML. Metastasis results from preexisting variant cells within a malignant tumor. *Science.* 1977 Aug 26;197(4306):893-5.
- Fidler IJ. Metastasis: Quantitative analysis of distribution and fate of tumor emboli labeled with 125 I-5-iodo-2'-deoxyuridine. *J Natl Cancer Inst.* 1970 Oct;45(4):773-82.
- Finnsen KW, Tam BY, Liu K, Marcoux A, Lepage P, Roy S, et al. Identification of CD109 as part of the TGF-beta receptor system in human keratinocytes. *FASEB J.* 2006 Jul;20(9):1525-7.
- Frehlick LJ, Eirin-Lopez JM, Ausio J. New insights into the nucleophosmin/nucleoplasmin family of nuclear chaperones. *Bioessays.* 2007 Jan;29(1):49-59.
- Friberg S, Mattson S. On the growth rates of human malignant tumors: Implications for medical decision making. *J Surg Oncol.* 1997 Aug;65(4):284-97.
- Fukawa T, Ono M, Matsuo T, Uehara H, Miki T, Nakamura Y, et al. DDX31 regulates the p53-HDM2 pathway and rRNA gene transcription through its interaction with NPM1 in renal cell carcinomas. *Cancer Res.* 2012 Nov 15;72(22):5867-77.
- Futreal PA. A census of human cancer genes. *Nature Reviews Cancer.* 2004;4(3):177-83.
- Gadad SS, Senapati P, Syed SH, Rajan RE, Shandilya J, Swaminathan V, et al. The multifunctional protein nucleophosmin (NPM1) is a human linker histone H1 chaperone. *Biochemistry.* 2011 Apr 12;50(14):2780-9.
- Gallo A, Lo Sterzo C, Mori M, Di Matteo A, Bertini I, Banci L, et al. Structure of nucleophosmin DNA-binding domain and analysis of its complex with a G-quadruplex sequence from the c-MYC promoter. *J Biol Chem.* 2012 Aug 3;287(32):26539-48.
- Gauthier ML, Berman HK, Miller C, Kozakeiwicz K, Chew K, Moore D, et al. Abrogated response to cellular stress identifies DCIS associated with subsequent tumor events and defines basal-like breast tumors. *Cancer Cell.* 2007 Nov;12(5):479-91.
- Ginestier C, Charafe-Jauffret E, Birnbaum D. What drives breast cancer heterogeneity: Oncogenic events or cell of origin? *J Pathol.* 2012 Jul;227(3):267-9.
- Goldhirsch A, Wood WC, Coates AS, Gelber RD, Thurlimann B, Senn HJ, et al. Strategies for subtypes--dealing with the diversity of breast cancer: Highlights of the st. gallen international expert consensus on the primary therapy of early breast cancer 2011. *Ann Oncol.* 2011 Aug;22(8):1736-47.
- Goldstein BJ, Ahmad F, Ding W, Li PM, Zhang WR. Regulation of the insulin signalling pathway by cellular protein-tyrosine phosphatases. *Mol Cell Biochem.* 1998 May;182(1-2):91-9.
- Goodison S, Kawai K, Hihara J, Jiang P, Yang M, Urquidí V, et al. Prolonged dormancy and site-specific growth potential of cancer cells spontaneously disseminated from nonmetastatic breast tumors as revealed by labeling with green fluorescent protein. *Clin Cancer Res.* 2003 Sep 1;9(10 Pt 1):3808-14.

Goss PE, Chambers AF. Does tumour dormancy offer a therapeutic target? *Nat Rev Cancer*. 2010 Dec;10(12):871-7.

Grisendi S, Mecucci C, Falini B, Pandolfi PP. Nucleophosmin and cancer. *Nat Rev Cancer*. 2006 Jul;6(7):493-505.

Grisendi S, Bernardi R, Rossi M, Cheng K, Khandker L, Manova K, et al. Role of nucleophosmin in embryonic development and tumorigenesis. *Nature*. 2005 Sep 1;437(7055):147-53.

Gritsenko PG, Ilina O, Friedl P. Interstitial guidance of cancer invasion. *J Pathol*. 2012 Jan;226(2):185-99.

Guo QM, Malek RL, Kim S, Chiao C, He M, Ruffey M, et al. Identification of c-myc responsive genes using rat cDNA microarray. *Cancer Res*. 2000 Nov 1;60(21):5922-8.

Gupta GP, Massague J. Cancer metastasis: Building a framework. *Cell*. 2006 Nov 17;127(4):679-95.

Hagiwara S, Murakumo Y, Mii S, Shigetomi T, Yamamoto N, Furue H, et al. Processing of CD109 by furin and its role in the regulation of TGF-beta signaling. *Oncogene*. 2010 Apr 15;29(15):2181-91.

Haindl M, Harasim T, Eick D, Muller S. The nucleolar SUMO-specific protease SENP3 reverses SUMO modification of nucleophosmin and is required for rRNA processing. *EMBO Rep*. 2008 Mar;9(3):273-9.

Hanahan D, Weinberg RA. Hallmarks of cancer: The next generation. *Cell*. 2011 Mar 4;144(5):646-74.

Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. 2000 Jan 7;100(1):57-70.

Hanahan D. Accessories to the crime: Functions of cells recruited to the tumor microenvironment. *Cancer Cell*. 2012;21(3):309-22.

Haneji T, Teramachi J, Hirashima K, Kimura K, Morimoto H. Interaction of protein phosphatase 1delta with nucleophosmin in human osteoblastic cells. *Acta Histochem Cytochem*. 2012 Feb 29;45(1):1-7.

Hart IR, Fidler IJ. Role of organ selectivity in the determination of metastatic patterns of B16 melanoma. *Cancer Res*. 1980 Jul;40(7):2281-7.

Hartman AR, Kaldate RR, Sailer LM, Painter L, Grier CE, Endsley RR, et al. Prevalence of BRCA mutations in an unselected population of triple-negative breast cancer. *Cancer*. 2012 Jun 1;118(11):2787-95.

Hasegawa M, Moritani S, Murakumo Y, Sato T, Hagiwara S, Suzuki C, et al. CD109 expression in basal-like breast carcinoma. *Pathol Int*. 2008 May;58(5):288-94.

Hedegaard M, Krafft C, Ditzel HJ, Johansen LE, Hassing S, Popp J. Discriminating isogenic cancer cells and identifying altered unsaturated fatty acid content as associated with metastasis status, using k-means clustering and partial least squares-discriminant analysis of raman maps. *Anal Chem*. 2010 Apr 1;82(7):2797-802.

Heiss MM, Allgayer H, Gruetzner KU, Funke I, Babic R, Jauch KW, et al. Individual development and uPA-receptor expression of disseminated tumour cells in bone marrow: A reference to early systemic disease in solid cancer. *Nat Med*. 1995 Oct;1(10):1035-9.

Herrera JE, Correia JJ, Jones AE, Olson MO. Sedimentation analyses of the salt- and divalent metal ion-induced oligomerization of nucleolar protein B23. *Biochemistry*. 1996 Feb 27;35(8):2668-73.

## REFERENCES

- Herrera JE, Savkur R, Olson MO. The ribonuclease activity of nucleolar protein B23. *Nucleic Acids Res.* 1995 Oct 11;23(19):3974-9.
- Herschkowitz JI, He X, Fan C, Perou CM. The functional loss of the retinoblastoma tumour suppressor is a common event in basal-like and luminal B breast carcinomas. *Breast Cancer Res.* 2008;10(5):R75.
- Hingorani K, Szebeni A, Olson MOJ. Mapping the functional domains of nucleolar protein B23. *J Biol Chem.* 2000;275(32):24451.
- Hisaoka M, Ueshima S, Murano K, Nagata K, Okuwaki M. Regulation of nucleolar chromatin by B23/nucleophosmin jointly depends upon its RNA binding activity and transcription factor UBF. *Mol Cell Biol.* 2010 Oct;30(20):4952-64.
- Hockla A, Radisky DC, Radisky ES. Mesotrypsin promotes malignant growth of breast cancer cells through shedding of CD109. *Breast Cancer Res Treat.* 2010 Nov;124(1):27-38.
- Hoffman B, Liebermann DA. Apoptotic signaling by c-MYC. *Oncogene.* 2008 Oct 27;27(50):6462-72.
- Hollestelle A, Nagel JH, Smid M, Lam S, Elstrodt F, Wasielewski M, et al. Distinct gene mutation profiles among luminal-type and basal-type breast cancer cell lines. *Breast Cancer Res Treat.* 2010 May;121(1):53-64.
- Hollestelle A, Schutte M. Comment re: MDA-MB-435 and M14 cell lines: Identical but not M14 melanoma? *Cancer Res.* 2009 Oct 1;69(19):7893,5472.CAN-09-2396. Epub 2009 Sep 1.
- Hollstein M, Sidransky D, Vogelstein B, Harris CC. P53 mutations in human cancers. *Science.* 1991 Jul 5;253(5015):49-53.
- Honda R, Yasuda H. Association of p19(ARF) with Mdm2 inhibits ubiquitin ligase activity of Mdm2 for tumor suppressor p53. *EMBO J.* 1999 Jan 4;18(1):22-7.
- Hondermarck H, Tastet C, El Yazidi-Belkoura I, Toillon RA, Le Bourhis X. Proteomics of breast cancer: The quest for markers and therapeutic targets. *J Proteome Res.* 2008 Apr;7(4):1403-11.
- Hopkins AL, Groom CR. The druggable genome. *Nat Rev Drug Discov.* 2002 Sep;1(9):727-30.
- Horak CE, Lee JH, Marshall JC, Shreeve SM, Steeg PS. The role of metastasis suppressor genes in metastatic dormancy. *APMIS.* 2008 Jul-Aug;116(7-8):586-601.
- Horn HF, Vousden KH. Cancer: Guarding the guardian? *Nature.* 2004 Jan 8;427(6970):110-1.
- Hsu CY, Yung BY. Down-regulation of nucleophosmin/B23 during retinoic acid-induced differentiation of human promyelocytic leukemia HL-60 cells. *Oncogene.* 1998 Feb 19;16(7):915-23.
- Hu Z, Fan C, Oh DS, Marron JS, He X, Qaqish BF, et al. The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics.* 2006 Apr 27;7:96.
- Huang H, Yee J, Shew J, Chen P, Bookstein R, Friedmann T, et al. Suppression of the neoplastic phenotype by replacement of the RB gene in human cancer cells. (retinoblastoma). *Science*, Dec 16, 1988, Vol 242(4885), p 1563(4). 1988;242(4885):1563.
- Hugh J, Hanson J, Cheang MC, Nielsen TO, Perou CM, Dumontet C, et al. Breast cancer subtypes and response to docetaxel in node-positive breast cancer: Use of an immunohistochemical definition in the BCIRG 001 trial. *J Clin Oncol.* 2009 Mar 10;27(8):1168-76.
- Hunter K, Welch DR, Liu ET. Genetic background is an important determinant of metastatic potential. *Nat Genet.* 2003 May;34(1):23,4; author reply 25.

Inouye CJ, Seto E. Relief of YY1-induced transcriptional repression by protein-protein interaction with the nucleolar phosphoprotein B23. *J Biol Chem.* 1994 Mar 4;269(9):6506-10.

Itahana K, Bhat KP, Jin A, Itahana Y, Hawke D, Kobayashi R, et al. Tumor suppressor ARF degrades B23, a nucleolar protein involved in ribosome biogenesis and cell proliferation. *Mol Cell.* 2003 Nov;12(5):1151-64.

Ji H, Ding Z, Hawke D, Xing D, Jiang BH, Mills GB, et al. AKT-dependent phosphorylation of riban regulates nucleophosmin- and MDM2-mediated p53 stability and cell apoptosis. *EMBO Rep.* 2012 Jun 1;13(6):554-60.

Jian Y, Gao Z, Sun J, Shen Q, Feng F, Jing Y, et al. RNA aptamers interfering with nucleophosmin oligomerization induce apoptosis of cancer cells. *Oncogene.* 2009 Nov 26;28(47):4201-11.

Joensuu H, Bono P, Kataja V, Alanko T, Kokko R, Asola R, et al. Fluorouracil, epirubicin, and cyclophosphamide with either docetaxel or vinorelbine, with or without trastuzumab, as adjuvant treatments of breast cancer: Final results of the FinHer trial. *Journal of Clinical Oncology.* 2009;27(34):5685.

Joensuu H, Kellokumpu-Lehtinen PL, Bono P, Alanko T, Kataja V, Asola R, et al. Adjuvant docetaxel or vinorelbine with or without trastuzumab for breast cancer. *N Engl J Med.* 2006 Feb 23;354(8):809-20.

Joensuu H, Isola J, Lundin M, Salminen T, Holli K, Kataja V, et al. Amplification of erbB2 and erbB2 expression are superior to estrogen receptor status as risk factors for distant recurrence in pT1N0M0 breast cancer: A nationwide population-based study. *Clin Cancer Res.* 2003 Mar;9(3):923-30.

Johnson RC, Augustin-Voss HG, Zhu DZ, Pauli BU. Endothelial cell membrane vesicles in the study of organ preference of metastasis. *Cancer Res.* 1991 Jan 1;51(1):394-9.

Joyce JA, Pollard JW. Microenvironmental regulation of metastasis. *Nat Rev Cancer.* 2009 Apr;9(4):239-52.

Junttila MR, Puustinen P, Niemela M, Ahola R, Arnold H, Bottzauw T, et al. CIP2A inhibits PP2A in human malignancies. *Cell.* 2007 Jul 13;130(1):51-62.

Kanehisa M, Goto S, Furumichi M, Tanabe M, Hirakawa M. KEGG for representation and analysis of molecular networks involving diseases and drugs. *Nucleic Acids Res.* 2010 Jan;38(Database issue):D355-60.

Kang YJ, Olson MO, Busch H. Phosphorylation of acid-soluble proteins in isolated nucleoli of novikoff hepatoma ascites cells. effects of divalent cations. *J Biol Chem.* 1974 Sep 10;249(17):5580-5.

Karhemo PR, Hyvonen M, Laakkonen P. Metastasis-associated cell surface oncoproteomics. *Front Pharmacol.* 2012;3:192.

Karrison TG, Ferguson DJ, Meier P. Dormancy of mammary carcinoma after mastectomy. *J Natl Cancer Inst.* 1999 Jan 6;91(1):80-5.

Khanna A, Okkeri J, Bilgen T, Tiirikka T, Vihinen M, Visakorpi T, et al. ETS1 mediates MEK1/2-dependent overexpression of cancerous inhibitor of protein phosphatase 2A (CIP2A) in human cancer cells. *PLoS One.* 2011 Mar 22;6(3):e17979.

Kikuta K, Tochigi N, Shimoda T, Yabe H, Morioka H, Toyama Y, et al. Nucleophosmin as a candidate prognostic biomarker of ewing's sarcoma revealed by proteomics. *Clin Cancer Res.* 2009 Apr 15;15(8):2885-94.

## REFERENCES

- Kim RS, Avivar-Valderas A, Estrada Y, Bragado P, Sosa MS, Aguirre-Ghiso JA, et al. Dormancy signatures and metastasis in estrogen receptor positive and negative breast cancer. *PLoS One*. 2012;7(4):e35569.
- Kim S, Li Q, Dang CV, Lee LA. Induction of ribosomal genes and hepatocyte hypertrophy by adenovirus-mediated expression of c-myc in vivo. *Proc Natl Acad Sci U S A*. 2000 Oct 10;97(21):11198-202.
- Klein CA. Parallel progression of primary tumours and metastases. *Nat Rev Cancer*. 2009 Apr;9(4):302-12.
- Knudson AG. GENETIC PREDISPOSITION TO CANCER. *Cancer Detect Prev*. 1984;7(1):1-8.
- Kononen J, Bubendorf L, Kallioniemi A, Bärklund M, Schraml P, Leighton S, et al. Tissue microarrays for high-throughput molecular profiling of tumor specimens. *Nat Med*. 1998;4(7):844.
- Korgaonkar C, Hagen J, Tompkins V, Frazier A, Allamargot C, Quelle F, et al. Nucleophosmin (B23) targets ARF to nucleoli and inhibits its function. *Molecular & Cellular Biology*. 2005;25(4):1258-71.
- Koscielny S, Tubiana M, Le MG, Valleron AJ, Mouriessé H, Contesso G, et al. Breast cancer: Relationship between the size of the primary tumour and the probability of metastatic dissemination. *Br J Cancer*. 1984 Jun;49(6):709-15.
- Krause A, Hoffmann I. Polo-like kinase 2-dependent phosphorylation of NPM/B23 on serine 4 triggers centriole duplication. *PLoS One*. 2010 Mar 24;5(3):e9849.
- Kuo ML, den Besten W, Bertwistle D, Roussel MF, Sherr CJ. N-terminal polyubiquitination and degradation of the arf tumor suppressor. *Genes Dev*. 2004 Aug 1;18(15):1862-74.
- Kurki S, Peltonen K, Latonen L, Kiviharju TM, Ojala PM, Meek D, et al. Nucleolar protein NPM interacts with HDM2 and protects tumor suppressor protein p53 from HDM2-mediated degradation. *Cancer Cell*. 2004 May;5(5):465-75.
- Kuukasjarvi T, Karhu R, Tanner M, Kahkonen M, Schaffer A, Nupponen N, et al. Genetic heterogeneity and clonal evolution underlying development of asynchronous metastasis in human breast cancer. *Cancer Res*. 1997 Apr 15;57(8):1597-604.
- Laakso M, Hautaniemi S. Integrative platform to translate gene sets to networks. *Bioinformatics*. 2010 Jul 15;26(14):1802-3.
- Laine A, Sihto H, Come C, Rosenfeldt MT, Zwolinska A, Niemela M, et al. Senescence sensitivity of breast cancer cells is defined by positive feedback loop between CIP2A and E2F1. *Cancer Discov*. 2013 Feb;3(2):182-97.
- Lam L, Aktary Z, Bishay M, Werkman C, Kuo C-, Heacock M, et al. Regulation of subcellular distribution and oncogenic potential of nucleophosmin by plakoglobin. *Oncogenesis*. 2012;1(3).
- Lambert B, Buckle M. Characterisation of the interface between nucleophosmin (NPM) and p53: Potential role in p53 stabilisation. *FEBS Lett*. 2006 Jan 9;580(1):345-50.
- Largillier R, Ferrero JM, Doyen J, Barriere J, Namer M, Mari V, et al. Prognostic factors in 1,038 women with metastatic breast cancer. *Ann Oncol*. 2008 Dec;19(12):2012-9.
- Lazo PA. Functional implications of tetraspanin proteins in cancer biology. *Cancer Sci*. 2007 Nov;98(11):1666-77.
- Lee HH, Kim HS, Kang JY, Lee BI, Ha JY, Yoon HJ, et al. Crystal structure of human nucleophosmin-core reveals plasticity of the pentamer-pentamer interface. *Proteins*. 2007 Nov 15;69(3):672-8.



- Lee SB, Xuan Nguyen TL, Choi JW, Lee KH, Cho SW, Liu Z, et al. Nuclear akt interacts with B23/NPM and protects it from proteolytic cleavage, enhancing cell survival. *Proc Natl Acad Sci U S A*. 2008 Oct 28;105(43):16584-9.
- Leotoing L, Meunier L, Manin M, Mauduit C, Decaussin M, Verrijdt G, et al. Influence of nucleophosmin/B23 on DNA binding and transcriptional activity of the androgen receptor in prostate cancer cell. *Oncogene*. 2008 May 1;27(20):2858-67.
- Leth-Larsen R, Lund RR, Ditzel HJ. Plasma membrane proteomics and its application in clinical cancer biomarker discovery. *Mol Cell Proteomics*. 2010 Jul;9(7):1369-82.
- Leth-Larsen R, Lund R, Hansen HV, Laenkholtm AV, Tarin D, Jensen ON, et al. Metastasis-related plasma membrane proteins of human breast cancer cells identified by comparative quantitative mass spectrometry. *Mol Cell Proteomics*. 2009 Jun;8(6):1436-49.
- Levine AJ, Oren M. The first 30 years of p53: Growing ever more complex. *Nat Rev Cancer*. 2009 Oct;9(10):749-58.
- Levine AJ, Momand J, Finlay CA. The p53 tumour suppressor gene. *Nature*. 1991 Jun 6;351(6326):453-6.
- Li J, Sejas DP, Burma S, Chen DJ, Pang Q. Nucleophosmin suppresses oncogene-induced apoptosis and senescence and enhances oncogenic cooperation in cells with genomic instability. *Carcinogenesis*. 2007 Jun;28(6):1163-70.
- Li J, Zhang X, Sejas DP, Bagby GC, Pang Q. Hypoxia-induced nucleophosmin protects cell death through inhibition of p53. *J Biol Chem*. 2004 Oct 1;279(40):41275-9.
- Li Z, Hann SR. Nucleophosmin is essential for c-myc nucleolar localization and c-myc-mediated rDNA transcription. *Oncogene*. 2013 Apr 11;32(15):1988-94.
- Li Z, Boone D, Hann SR. Nucleophosmin interacts directly with c-myc and controls c-myc-induced hyperproliferation and transformation. *Proc Natl Acad Sci U S A*. 2008 Dec 2;105(48):18794-9.
- Lin CY, Tan BCM, Liu H, Shih CJ, Chien KY, Lin CL, et al. Dephosphorylation of nucleophosmin by PP1 $\beta$  facilitates pRB binding and consequent E2F1-dependent DNA repair. *Mol Biol Cell*. 2010;21(24):4409.
- Lin CY, Liang YC, Yung BY. Nucleophosmin/B23 regulates transcriptional activation of E2F1 via modulating the promoter binding of NF-kappaB, E2F1 and pRB. *Cell Signal*. 2006 Nov;18(11):2041-8.
- Lipscomb EA, Mercurio AM. Mobilization and activation of a signaling competent alpha6beta4integrin underlies its contribution to carcinoma progression. *Cancer Metastasis Rev*. 2005 Sep;24(3):413-23.
- Liu D, Aguirre Ghiso J, Estrada Y, Ossowski L. EGFR is a transducer of the urokinase receptor initiated signal that is required for in vivo growth of a human carcinoma. *Cancer Cell*. 2002 Jun;1(5):445-57.
- Liu H, Tan BC, Tseng KH, Chuang CP, Yeh CW, Chen KD, et al. Nucleophosmin acts as a novel AP2alpha-binding transcriptional corepressor during cell differentiation. *EMBO Rep*. 2007 Apr;8(4):394-400.
- Liu QR, Chan PK. Formation of nucleophosmin/B23 oligomers requires both the amino- and the carboxyl-terminal domains of the protein. *Eur J Biochem*. 1991 Sep 15;200(3):715-21.
- Liu S, Goldstein RH, Scepansky EM, Rosenblatt M. Inhibition of rho-associated kinase signaling prevents breast cancer metastasis to human bone. *Cancer Res*. 2009 Nov 15;69(22):8742-51.

## REFERENCES

- Liu X, Liu Z, Jang SW, Ma Z, Shinmura K, Kang S, et al. Sumoylation of nucleophosmin/B23 regulates its subcellular localization, mediating cell proliferation and survival. *Proc Natl Acad Sci U S A*. 2007 Jun 5;104(23):9679-84.
- Liu Y, Zhang F, Zhang XF, Qi LS, Yang L, Guo H, et al. Expression of nucleophosmin/NPM1 correlates with migration and invasiveness of colon cancer cells. *J Biomed Sci*. 2012 May 25;19:53,0127-19-53.
- Lorusso G, Ruegg C. New insights into the mechanisms of organ-specific breast cancer metastasis. *Semin Cancer Biol*. 2012 Jun;22(3):226-33.
- Lu P, Takai K, Weaver VM, Werb Z. Extracellular matrix degradation and remodeling in development and disease. *Cold Spring Harb Perspect Biol*. 2011 Dec 1;3(12):10.1101/cshperspect.a005058.
- Lu X, Kang Y. Organotropism of breast cancer metastasis. *J Mammary Gland Biol Neoplasia*. 2007 Sep;12(2-3):153-62.
- Lund R, Leth-Larsen R, Jensen ON, Ditzel HJ. Efficient isolation and quantitative proteomic analysis of cancer cell plasma membrane proteins for identification of metastasis-associated cell surface markers. *J Proteome Res*. 2009 Jun;8(6):3078-90.
- Luque-Garcia JL, Martinez-Torrecuadrada JL, Epifano C, Canamero M, Babel I, Casal JJ. Differential protein expression on the cell surface of colorectal cancer cells associated to tumor metastasis. *Proteomics*. 2010 Mar;10(5):940-52.
- Ma Z, Kanai M, Kawamura K, Kaibuchi K, Ye K, Fukasawa K. Interaction between ROCK II and nucleophosmin/B23 in the regulation of centrosome duplication. *Mol Cell Biol*. 2006 Dec;26(23):9016-34.
- Maggi LB, Jr, Kuchenruether M, Dadey DY, Schwoppe RM, Grisendi S, Townsend RR, et al. Nucleophosmin serves as a rate-limiting nuclear export chaperone for the mammalian ribosome. *Mol Cell Biol*. 2008 Dec;28(23):7050-65.
- Maiguel DA, Jones L, Chakravarty D, Yang C, Carrier F. Nucleophosmin sets a threshold for p53 response to UV radiation. *Mol Cell Biol*. 2004 May;24(9):3703-11.
- Markey MP, Bergseid, J.,: Bosco, E., Stengel K, Xu H, Mayhew C, N., Schwemberger S, J., et al. Loss of the retinoblastoma tumor suppressor: Differential action on transcriptional programs related to cell cycle control and immune function.(original article)(clinical report). *Oncogene*, Sept 20, 2007, Vol 26(43), p 6307. 2007;26(43):6307.
- Mehlen P, Puisieux A. Metastasis: A question of life or death. *Nat Rev Cancer*. 2006 Jun;6(6):449-58.
- Mendes-da-Silva P, Moreira A, Duro-da-Costa J, Matias D, Monteiro C. Frequent loss of heterozygosity on chromosome 5 in non-small cell lung carcinoma. *Mol Pathol*. 2000 Aug;53(4):184-7.
- Meng S, Tripathy D, Frenkel EP, Shete S, Naftalis EZ, Huth JF, et al. Circulating tumor cells in patients with breast cancer dormancy. *Clin Cancer Res*. 2004 Dec 15;10(24):8152-62.
- Mitre DM, Kriwacki RW. Cryptic disorder: An order-disorder transformation regulates the function of nucleophosmin. *Pac Symp Biocomput*. 2012:152-63.
- Mittnacht S. Control of pRB phosphorylation. *Curr Opin Genet Dev*. 1998;8(1):21-7.
- Molyneux G, Smalley MJ. The cell of origin of BRCA1 mutation-associated breast cancer: A cautionary tale of gene expression profiling. *J Mammary Gland Biol Neoplasia*. 2011 Apr;16(1):51-5.

Montel V, Suzuki M, Galloy C, Mose ES, Tarin D. Expression of melanocyte-related genes in human breast cancer and its implications. *Differentiation*. 2009 Dec;78(5):283-91.

Montel V, Mose ES, Tarin D. Tumor-stromal interactions reciprocally modulate gene expression patterns during carcinogenesis and metastasis. *Int J Cancer*. 2006 Jul 15;119(2):251-63.

Montel V, Huang TY, Mose E, Pestonjamas K, Tarin D. Expression profiling of primary tumors and matched lymphatic and lung metastases in a xenogeneic breast cancer model. *Am J Pathol*. 2005 May;166(5):1565-79.

Muller A, Homey B, Soto H, Ge N, Catron D, Buchanan ME, et al. Involvement of chemokine receptors in breast cancer metastasis. *Nature*. 2001 Mar 1;410(6824):50-6.

Muller V, Stahmann N, Riethdorf S, Rau T, Zabel T, Goetz A, et al. Circulating tumor cells in breast cancer: Correlation to bone marrow micrometastases, heterogeneous response to systemic therapy and low proliferative activity. *Clin Cancer Res*. 2005 May 15;11(10):3678-85.

Muralidharan-Chari V, Clancy JW, Sedgwick A, D'Souza-Schorey C. Microvesicles: Mediators of extracellular communication during cancer progression. *J Cell Sci*. 2010 May 15;123(Pt 10):1603-11.

Murano K, Okuwaki M, Hisaoka M, Nagata K. Transcription regulation of the rRNA gene by a multifunctional nucleolar protein, B23/nucleophosmin, through its histone chaperone activity. *Mol Cell Biol*. 2008 May;28(10):3114-26.

Nalabothula N, Chakravarty D, Pierce A, Carrier F. Over expression of nucleophosmin and nucleolin contributes to the suboptimal activation of a G2/M checkpoint in ataxia telangiectasia fibroblasts. *Mol Cell Pharmacol*. 2010;2(5):179-89.

Naumov GN, Bender E, Zurakowski D, Kang SY, Sampson D, Flynn E, et al. A model of human tumor dormancy: An angiogenic switch from the nonangiogenic phenotype. *J Natl Cancer Inst*. 2006 Mar 1;98(5):316-25.

Naumov GN, MacDonald IC, Weinmeister PM, Kerkvliet N, Nadkarni KV, Wilson SM, et al. Persistence of solitary mammary carcinoma cells in a secondary site: A possible contributor to dormancy. *Cancer Res*. 2002 Apr 1;62(7):2162-8.

Negi SS, Olson MO. Effects of interphase and mitotic phosphorylation on the mobility and location of nucleolar protein B23. *J Cell Sci*. 2006 Sep 1;119(Pt 17):3676-85.

Neiman PE, Ruddell A, Jasoni C, Loring G, Thomas SJ, Brandvold KA, et al. Analysis of gene expression during myc oncogene-induced lymphomagenesis in the bursa of fabricius. *Proc Natl Acad Sci U S A*. 2001 May 22;98(11):6378-83.

Nerlich AG, Bachmeier BE. Density-dependent lineage instability of MDA-MB-435 breast cancer cells. *Oncol Lett*. 2013 Apr;5(4):1370-4.

Neve RM, Chin K, Fridlyand J, Yeh J, Baehner FL, Fevr T, et al. A collection of breast cancer cell lines for the study of functionally distinct cancer subtypes. *Cancer Cell*. 2006 Dec;10(6):515-27.

Nguyen DX, Bos PD, Massague J. Metastasis: From dissemination to organ-specific colonization. *Nat Rev Cancer*. 2009 Apr;9(4):274-84.

Nielsen TO, Hsu FD, Jensen K, Cheang M, Karaca G, Hu Z, et al. Immunohistochemical and clinical characterization of the basal-like subtype of invasive breast carcinoma. *Clin Cancer Res*. 2004 Aug 15;10(16):5367-74.

## REFERENCES

- Niemela M, Kauko O, Sihto H, Mpindi JP, Nicorici D, Pernila P, et al. CIP2A signature reveals the MYC dependency of CIP2A-regulated phenotypes and its clinical association with breast cancer subtypes. *Oncogene*. 2012 Sep 27;31(39):4266-78.
- Nilsson JA, Cleveland JL. Myc pathways provoking cell suicide and cancer. *Oncogene*. 2003 Dec 8;22(56):9007-21.
- Nishimura Y, Ohkubo T, Furuichi Y, Umekawa H. Tryptophans 286 and 288 in the C-terminal region of protein B23.1 are important for its nucleolar localization. *Biosci Biotechnol Biochem*. 2002 Oct;66(10):2239-42.
- Norton L, Massague J. Is cancer a disease of self-seeding? *Nat Med*. 2006 Aug;12(8):875-8.
- Nozawa Y, Van Belzen N, Van der Made AC, Dinjens WN, Bosman FT. Expression of nucleophosmin/B23 in normal and neoplastic colorectal mucosa. *J Pathol*. 1996 Jan;178(1):48-52.
- Ochs R, Lischwe M, O'Leary P, Busch H. Localization of nucleolar phosphoproteins B23 and C23 during mitosis. *Exp Cell Res*. 1983 Jun;146(1):139-49.
- O'Connor PM, Jackman J, Bae I, Myers TG, Fan S, Mutoh M, et al. Characterization of the p53 tumor suppressor pathway in cell lines of the national cancer institute anticancer drug screen and correlations with the growth-inhibitory potency of 123 anticancer agents. *Cancer Res*. 1997 Oct 1;57(19):4285-300.
- Ohshima Y, Yajima I, Kumasaka MY, Yanagishita T, Watanabe D, Takahashi M, et al. CD109 expression levels in malignant melanoma. *J Dermatol Sci*. 2010 Feb;57(2):140-2.
- Okuda M. The role of nucleophosmin in centrosome duplication. *Oncogene*. 2002 Sep 9;21(40):6170-4.
- Okuda M, Horn HF, Tarapore P, Tokuyama Y, Smulian AG, Chan PK, et al. Nucleophosmin/B23 is a target of CDK2/cyclin E in centrosome duplication. *Cell*. 2000 Sep 29;103(1):127-40.
- Okuwaki M, Tsujimoto M, Nagata K. The RNA binding activity of a ribosome biogenesis factor, nucleophosmin/B23, is modulated by phosphorylation with a cell cycle-dependent kinase and by association with its subtype. *Mol Biol Cell*. 2002;13(6):2016.
- Okuwaki M, Matsumoto K, Tsujimoto M, Nagata K. Function of nucleophosmin/B23, a nucleolar acidic protein, as a histone chaperone. *FEBS Lett*. 2001;506(3):272.
- Ong SE, Mann M. Mass spectrometry-based proteomics turns quantitative. *Nat Chem Biol*. 2005 Oct;1(5):252-62.
- Orrick LR, Olson MO, Busch H. Comparison of nucleolar proteins of normal rat liver and novikoff hepatoma ascites cells by two-dimensional polyacrylamide gel electrophoresis. *Proc Natl Acad Sci U S A*. 1973 May;70(5):1316-20.
- Overington JP, Al-Lazikani B, Hopkins AL. How many drug targets are there? *Nat Rev Drug Discov*. 2006 Dec;5(12):993-6.
- Paget S. The distribution of secondary growths in cancer of the breast. *Lancet*. 1889;133(3421):571-3.
- Palmieri D, Smith QR, Lockman PR, Bronder J, Gril B, Chambers AF, et al. Brain metastases of breast cancer. *Breast Dis*. 2006 -2007;26:139-47.
- Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature*. 2000 Aug 17;406(6797):747-52.

Peter M, Nakagawa J, Doree M, Labbe JC, Nigg EA. Identification of major nucleolar proteins as candidate mitotic substrates of cdc2 kinase. *Cell*. 1990;60(5):791.

Place AE, Jin Huh S, Polyak K. The microenvironment in breast cancer progression: Biology and implications for treatment. *Breast Cancer Res*. 2011;13(6):227.

Pontier SM, Muller WJ. Integrins in breast cancer dormancy. *APMIS*. 2008 Jul-Aug;116(7-8):677-84.

Poste G, Fidler IJ. The pathogenesis of cancer metastasis. *Nature*. 1980 Jan 10;283(5743):139-46.

Prat A, Parker JS, Karginova O, Fan C, Livasy C, Herschkowitz JI, et al. Phenotypic and molecular characterization of the claudin-low intrinsic subtype of breast cancer. *Breast Cancer Res*. 2010;12(5):R68.

Prestayko AW, Olson MO, Busch H. Phosphorylation of proteins of ribosomes and nucleolar preribosomal particles in vivo in novikoff hepatoma ascites cells. *FEBS Lett*. 1974 Aug 25;44(2):131-5.

Prinos P, Lacoste MC, Wong J, Bonneau AM, Georges E. Mutation of cysteine 21 inhibits nucleophosmin/B23 oligomerization and chaperone activity. *Int J Biochem Mol Biol*. 2011;2(1):24-30.

Psaila B, Lyden D. The metastatic niche: Adapting the foreign soil. *Nat Rev Cancer*. 2009 Apr;9(4):285-93.

Psaila B, Kaplan RN, Port ER, Lyden D. Priming the 'soil' for breast cancer metastasis: The pre-metastatic niche. *Breast Dis*. 2006 -2007;26:65-74.

Qi W, Shakalya K, Stejskal A, Goldman A, Beeck S, Cooke L, et al. NSC348884, a nucleophosmin inhibitor disrupts oligomer formation and induces apoptosis in human cancer cells. *Oncogene*. 2008 Jul 10;27(30):4210-20.

Qin Y, Hurley LH. Structures, folding patterns, and functions of intramolecular DNA G-quadruplexes found in eukaryotic promoter regions. *Biochimie*. 2008 Aug;90(8):1149-71.

Rae JM, Creighton CJ, Meck JM, Haddad BR, Johnson MD. MDA-MB-435 cells are derived from M14 melanoma cells--a loss for breast cancer, but a boon for melanoma research. *Breast Cancer Res Treat*. 2007 Jul;104(1):13-9.

Rae JM, Ramus SJ, Waltham M, Armes JE, Campbell IG, Clarke R, et al. Common origins of MDA-MB-435 cells from various sources with those shown to have melanoma properties. *Clin Exp Metastasis*. 2004;21(6):543-52.

Ramaswamy S, Ross KN, Lander ES, Golub TR. A molecular signature of metastasis in primary solid tumors. *Nat Genet*. 2003 Jan;33(1):49-54.

Ramos-Echazabal G, Chinea G, Garcia-Fernandez R, Pons T. In silico studies of potential phosphoresidues in the human nucleophosmin/B23: Its kinases and related biological processes. *J Cell Biochem*. 2012 Jul;113(7):2364-74.

Ramsamooj P, Notario V, Dritschilo A. Modification of nucleolar protein B23 after exposure to ionizing radiation. *Radiat Res*. 1995;143(2):158.

Rasmussen N, Ditzel HJ. Scanning the cell surface proteome of cancer cells and identification of metastasis-associated proteins using a subtractive immunization strategy. *J Proteome Res*. 2009 Nov;8(11):5048-59.

Recamier J. <br />Recherches sur le traitement du cancer par la compression méthodique simple ou combinée, et sur l'histoire générale de la même maladie. 2nd ed. ed. Paris: Gabon; 1829.

## REFERENCES

- Reis-Filho JS, Tutt AN. Triple negative tumours: A critical review. *Histopathology*. 2008 Jan;52(1):108-18.
- Reya T. Stem cells, cancer, and cancer stem cells. *Nature*, Nov 1, 2001, Vol 414(6859), p 105(7). 2001;414(6859):105.
- Roesli C, Borgia B, Schliemann C, Gunthert M, Wunderli-Allenspach H, Giavazzi R, et al. Comparative analysis of the membrane proteome of closely related metastatic and nonmetastatic tumor cells. *Cancer Res*. 2009 Jul 1;69(13):5406-14.
- Roesli C, Borgia B, Schliemann C, Gunthert M, Wunderli-Allenspach H, Giavazzi R, et al. Comparative analysis of the membrane proteome of closely related metastatic and nonmetastatic tumor cells. *Cancer Res*. 2009 Jul 1;69(13):5406-14.
- Roesli C, Mumprecht V, Neri D, Detmar M. Identification of the surface-accessible, lineage-specific vascular proteome by two-dimensional peptide mapping. *FASEB J*. 2008 Jun;22(6):1933-44.
- Ross DT, Scherf U, Eisen MB, Perou CM, Rees C, Spellman P, et al. Systematic variation in gene expression patterns in human cancer cell lines. *Nat Genet*. 2000 Mar;24(3):227-35.
- Ross JS, Slodkowska EA. Circulating and disseminated tumor cells in the management of breast cancer. *Am J Clin Pathol*. 2009 Aug;132(2):237-45.
- Rower C, Koy C, Hecker M, Reimer T, Gerber B, Thiesen HJ, et al. Mass spectrometric characterization of protein structure details refines the proteome signature for invasive ductal breast carcinoma. *J Am Soc Mass Spectrom*. 2011 Mar;22(3):440-56.
- Rower C, Ziems B, Radtke A, Schmitt O, Reimer T, Koy C, et al. Toponostics of invasive ductal breast carcinoma: Combination of spatial protein expression imaging and quantitative proteome signature analysis. *Int J Clin Exp Pathol*. 2011 Mar 31;4(5):454-67.
- Santos SC, Cavalli IJ, Ribeiro EM, Urban CA, Lima RS, Bleggi-Torres LF, et al. Patterns of DNA copy number changes in sentinel lymph node breast cancer metastases. *Cytogenet Genome Res*. 2008;122(1):16-21.
- Sarek G, Jarviluoma A, Moore HM, Tojkander S, Vartia S, Biberfeld P, et al. Nucleophosmin phosphorylation by v-cyclin-CDK6 controls KSHV latency. *PLoS Pathog*. 2010 Mar 19;6(3):e1000818.
- Sastry SK, Elferink LA. Checks and balances: Interplay of RTKs and PTPs in cancer progression. *Biochem Pharmacol*. 2011 Sep 1;82(5):435-40.
- Sato K, Hayami R, Wu W, Nishikawa T, Nishikawa H, Okuda Y, et al. Nucleophosmin/B23 is a candidate substrate for the BRCA1-BARD1 ubiquitin ligase. *J Biol Chem*. 2004 Jul 23;279(30):30919-22.
- Savkur RS, Olson MO. Preferential cleavage in pre-ribosomal RNA by protein B23 endoribonuclease. *Nucleic Acids Res*. 1998 Oct 1;26(19):4508-15.
- Schardt JA, Meyer M, Hartmann CH, Schubert F, Schmidt-Kittler O, Fuhrmann C, et al. Genomic analysis of single cytokeratin-positive cells from bone marrow reveals early mutational events in breast cancer. *Cancer Cell*. 2005 Sep;8(3):227-39.
- Scheurer SB, Rybak JN, Roesli C, Brunisholz RA, Potthast F, Schlapbach R, et al. Identification and relative quantification of membrane proteins by surface biotinylation and two-dimensional peptide mapping. *Proteomics*. 2005 Jul;5(11):2718-28.

Schmidt-Kittler O, Ragg T, Daskalakis A, Granzow M, Ahr A, Blankenstein TJ, et al. From latent disseminated cells to overt metastasis: Genetic analysis of systemic breast cancer progression. *Proc Natl Acad Sci U S A*. 2003 Jun 24;100(13):7737-42.

Schmidt-Zachmann MS, Hugle-Dorr B, Franke WW. A constitutive nucleolar protein identified as a member of the nucleoplasmin family. *EMBO J*. 1987 Jul;6(7):1881-90.

Schopper D, de Wolf C. How effective are breast cancer screening programmes by mammography? review of the current evidence. *Eur J Cancer*. 2009 Jul;45(11):1916-23.

Sellappan S, Grijalva R, Zhou X, Yang W, Eli MB, Mills GB, et al. Lineage infidelity of MDA-MB-435 cells: Expression of melanocyte proteins in a breast cancer cell line. *Cancer Res*. 2004 May 15;64(10):3479-85.

Shandilya J, Swaminathan V, Gadad SS, Choudhari R, Kodaganur GS, Kundu TK. Acetylated NPM1 localizes in the nucleoplasm and regulates transcriptional activation of genes implicated in oral cancer manifestation. *Mol Cell Biol*. 2009 Sep;29(18):5115-27.

Shibue T, Weinberg RA. Integrin beta1-focal adhesion kinase signaling directs the proliferation of metastatic cancer cells disseminated in the lungs. *Proc Natl Acad Sci U S A*. 2009 Jun 23;106(25):10290-5.

Shinmura K, Tarapore P, Tokuyama Y, George KR, Fukasawa K. Characterization of centrosomal association of nucleophosmin/B23 linked to Crml activity. *FEBS Lett*. 2005 Dec 5;579(29):6621-34.

Siddiqui-Jain A, Grand CL, Bearss DJ, Hurley LH. Direct evidence for a G-quadruplex in a promoter region and its targeting with a small molecule to repress c-MYC transcription. *Proc Natl Acad Sci U S A*. 2002 Sep 3;99(18):11593-8.

Sihto H, Lundin J, Lundin M, Lehtimäki T, Ristimäki A, Holli K, et al. Breast cancer biological subtypes and protein expression predict for the preferential distant metastasis sites: A nationwide cohort study. *Breast Cancer Res*. 2011 Sep 13;13(5):R87.

Skaar TC, Prasad SC, Sharareh S, Lippman ME, Brunner N, Clarke R. Two-dimensional gel electrophoresis analyses identify nucleophosmin as an estrogen regulated protein associated with acquired estrogen-independence in human breast cancer cells. *J Steroid Biochem Mol Biol*. 1998 Dec;67(5-6):391-402.

Sleeman JP, Christofori G, Fodde R, Collard JG, Berx G, Decraene C, et al. Concepts of metastasis in flux: The stromal progression model. *Semin Cancer Biol*. 2012 Jun;22(3):174-86.

Sleeman JP, Cremers N. New concepts in breast cancer metastasis: Tumor initiating cells and the microenvironment. *Clin Exp Metastasis*. 2007;24(8):707-15.

Smid M, Wang Y, Zhang Y, Sieuwerts AM, Yu J, Klijn JG, et al. Subtypes of breast cancer show preferential site of relapse. *Cancer Res*. 2008 May 1;68(9):3108-14.

So CH, Michal AM, Mashayekhi R, Benovic JL. G protein-coupled receptor kinase 5 phosphorylates nucleophosmin and regulates cell sensitivity to polo-like kinase 1 inhibition. *J Biol Chem*. 2012 May 18;287(21):17088-99.

Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, et al. Repeated observation of breast tumor subtypes in independent gene expression data sets. *Proc Natl Acad Sci U S A*. 2003 Jul 8;100(14):8418-23.

Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A*. 2001 Sep 11;98(19):10869-74.

## REFERENCES

- Spector DL, Ochs RL, Busch H. Silver staining, immunofluorescence, and immunoelectron microscopic localization of nucleolar phosphoproteins B23 and C23. *Chromosoma*. 1984;90(2):139-48.
- Sportoletti P, Grisendi S, Majid SM, Cheng K, Clohessy JG, Viale A, et al. Npm1 is a haploinsufficient suppressor of myeloid and lymphoid malignancies in the mouse. *Blood*. 2008 Apr 1;111(7):3859-62.
- Srinivasan S, Mahowald AP, Fuller MT. The receptor tyrosine phosphatase lar regulates adhesion between drosophila male germline stem cells and the niche. *Development*. 2012 Apr;139(8):1381-90.
- Steeg PS. New insights into the tumor metastatic process revealed by gene expression profiling. *Am J Pathol*. 2005 May;166(5):1291-4.
- Subhawong AP, Subhawong T, Nassar H, Kouprina N, Begum S, Vang R, et al. Most basal-like breast carcinomas demonstrate the same rb-/p16+ immunophenotype as the HPV-related poorly differentiated squamous cell carcinomas which they resemble morphologically. *Am J Surg Pathol*. 2009 Feb;33(2):163-75.
- Sugimoto M, Kuo ML, Roussel MF, Sherr CJ. Nucleolar arf tumor suppressor inhibits ribosomal RNA processing. *Mol Cell*. 2003 Feb;11(2):415-24.
- Suzuki M, Mose ES, Montel V, Tarin D. Dormant cancer cells retrieved from metastasis-free organs regain tumorigenic and metastatic potency. *Am J Pathol*. 2006 Aug;169(2):673-81.
- Swaminathan V, Kishore AH, Febitha KK, Kundu TK. Human histone chaperone nucleophosmin enhances acetylation-dependent chromatin transcription. *Mol Cell Biol*. 2005 Sep;25(17):7534-45.
- Szebeni A, Olson MOJ. Nucleolar protein B23 has molecular chaperone activities. *Protein Science*. 1999;8(4):905.
- Tago K, Chiocca S, Sherr CJ. Sumoylation induced by the arf tumor suppressor: A p53-independent function. *Proc Natl Acad Sci U S A*. 2005 May 24;102(21):7689-94.
- Takemura M, Ohoka F, Perpelescu M, Ogawa M, Matsushita H, Takaba T, et al. Phosphorylation-dependent migration of retinoblastoma protein into the nucleolus triggered by binding to nucleophosmin/B23. *Exp Cell Res*. 2002;276(2):233.
- Takemura M, Sato K, Nishio M, Akiyama T, Umekawa H, Yoshida S. Nucleolar protein B23.1 binds to retinoblastoma protein and synergistically stimulates DNA polymerase  $\alpha$ ; Activity1. *J Biochem*. 1999;125(5):904-9.
- Tarapore P, Okuda M, Fukasawa K. A mammalian in vitro centriole duplication system: Evidence for involvement of CDK2/cyclin E and nucleophosmin/B23 in centrosome duplication. *Cell Cycle*. 2002 Jan;1(1):75-81.
- Tokuyama Y, Horn HF, Kawamura K, Tarapore P, Fukasawa K. Specific phosphorylation of nucleophosmin on thr(199) by cyclin-dependent kinase 2-cyclin E and its role in centrosome duplication. *J Biol Chem*. 2001 Jun 15;276(24):21529-37.
- Torres L, Ribeiro FR, Pandis N, Andersen JA, Heim S, Teixeira MR. Intratumor genomic heterogeneity in breast cancer with clonal divergence between primary carcinomas and lymph node metastases. *Breast Cancer Res Treat*. 2007 Apr;102(2):143-55.
- Tsubura A, Yuri T, Yoshizawa K, Uehara N, Takada H. Role of fatty acids in malignancy and visual impairment: Epidemiological evidence and experimental studies. *Histol Histopathol*. 2009 Feb;24(2):223-34.



- Tsui KH, Juang HH, Lee TH, Chang PL, Chen CL, Yung BY. Association of nucleophosmin/B23 with bladder cancer recurrence based on immunohistochemical assessment in clinical samples. *Acta Pharmacol Sin*. 2008 Mar;29(3):364-70.
- Turner NC, Reis-Filho JS. Basal-like breast cancer and the BRCA1 phenotype. *Oncogene*. 2006 Sep 25;25(43):5846-53.
- Uhr JW, Pantel K. Controversies in clinical cancer dormancy. *Proc Natl Acad Sci U S A*. 2011 Jul 26;108(30):12396-400.
- Umekawa H, Chang JH, Correia JJ, Wang D, Wingfield PT, Olson MO. Nucleolar protein B23: Bacterial expression, purification, oligomerization and secondary structures of two isoforms. *Cell Mol Biol Res*. 1993;39(7):635-45.
- Urquidi V, Sloan D, Kawai K, Agarwal D, Woodman AC, Tarin D, et al. Contrasting expression of thrombospondin-1 and osteopontin correlates with absence or presence of metastatic phenotype in an isogenic model of spontaneous human breast cancer metastasis. *Clin Cancer Res*. 2002 Jan;8(1):61-74.
- Valastyan S, Weinberg RA. Tumor metastasis: Molecular insights and evolving paradigms. *Cell*. 2011 Oct 14;147(2):275-92.
- Vidwans SJ, Flaherty KT, Fisher DE, Tenenbaum JM, Travers MD, Shrager J. A melanoma molecular disease model. *PLoS ONE*. 2011;6(3).
- Voduc KD, Cheang MC, Tyldesley S, Gelmon K, Nielsen TO, Kennecke H. Breast cancer subtypes and the risk of local and regional relapse. *J Clin Oncol*. 2010 Apr 1;28(10):1684-91.
- Vydra J, Selicharova I, Smutna K, Sanda M, Matouskova E, Bursikova E, et al. Two-dimensional electrophoretic comparison of metastatic and non-metastatic human breast tumors using in vitro cultured epithelial cells derived from the cancer tissues. *BMC Cancer*. 2008 Apr 16;8:107,2407-8-107.
- Wang D, Baumann A, Szebeni A, Olson MOJ. The nucleic acid binding activity of nucleolar protein B23.1 resides in its carboxyl-terminal end. *J Biol Chem*. 1994;269(49):30994.
- Wang D, Umekawa H, Olson MO. Expression and subcellular locations of two forms of nucleolar protein B23 in rat tissues and cells. *Cell Mol Biol Res*. 1993;39(1):33-42.
- Wang J, Loberg R, Taichman RS. The pivotal role of CXCL12 (SDF-1)/CXCR4 axis in bone metastasis. *Cancer Metastasis Rev*. 2006 Dec;25(4):573-87.
- Wang Q, Hirohashi Y, Furuuchi K, Zhao H, Liu Q, Zhang H, et al. The centrosome in normal and transformed cells. *DNA Cell Biol*. 2004 Aug;23(8):475-89.
- Wang W, Budhu A, Forgues M, Wang XW. Temporal and spatial control of nucleophosmin by the ran-Crm1 complex in centrosome duplication. *Nat Cell Biol*. 2005 Aug;7(8):823-30.
- Weigelt B, Peterse JL, van 't Veer LJ. Breast cancer metastasis: Markers and models. *Nat Rev Cancer*. 2005 Aug;5(8):591-602.
- West L, Vidwans SJ, Campbell NP, Shrager J, Simon GR, Bueno R, et al. A novel classification of lung cancer into molecular subtypes. *PLoS ONE*. 2012;7(2).
- Willis RA. The spread of tumors in the human body. London: Churchill; 1934.
- Wojtukiewicz MZ, Zacharski LR, Memoli VA, Kisiel W, Kudryk BJ, Rousseau SM, et al. Malignant melanoma. interaction with coagulation and fibrinolysis pathways in situ. *Am J Clin Pathol*. 1990 Apr;93(4):516-21.
- Wozniak MA, Modzelewska K, Kwong L, Keely PJ. Focal adhesion regulation of cell behavior. *Biochim Biophys Acta*. 2004 Jul 5;1692(2-3):103-19.

## REFERENCES

- Wu MH, Chang JH, Yung BY. Resistance to UV-induced cell-killing in nucleophosmin/B23 over-expressed NIH 3T3 fibroblasts: Enhancement of DNA repair and up-regulation of PCNA in association with nucleophosmin/B23 over-expression. *Carcinogenesis*. 2002 Jan;23(1):93-100.
- Wu MH, Yung BY. UV stimulation of nucleophosmin/B23 expression is an immediate-early gene response induced by damaged DNA. *J Biol Chem*. 2002 Dec 13;277(50):48234-40.
- Xu Y, Fang F, Dhar SK, St Clair WH, Kasarskis EJ, St Clair DK. The role of a single-stranded nucleotide loop in transcriptional regulation of the human *sod2* gene. *J Biol Chem*. 2007 Jun 1;282(22):15981-94.
- Yang C, Maignel DA, Carrier F. Identification of nucleolin and nucleophosmin as genotoxic stress-responsive RNA-binding proteins. *Nucleic Acids Res*. 2002 May 15;30(10):2251-60.
- Yao J, Fu C, Ding X, Guo Z, Zenreski A, Chen Y, et al. Nek2A kinase regulates the localization of numatrin to centrosome in mitosis. *FEBS Lett*. 2004 Sep 24;575(1-3):112-8.
- You BJ, Huang IJ, Liu WH, Hung YB, Chang JH, Yung BY. Decrease in nucleophosmin/B23 mRNA and telomerase activity during indomethacin-induced apoptosis of gastric KATO-III cancer cells. *Naunyn Schmiedebergs Arch Pharmacol*. 1999 Dec;360(6):683-90.
- Yu Y, Maggi LB, Jr, Brady SN, Apicelli AJ, Dai MS, Lu H, et al. Nucleophosmin is essential for ribosomal protein L5 nuclear export. *Mol Cell Biol*. 2006 May;26(10):3798-809.
- Yun C, Wang Y, Mukhopadhyay D, Backlund P, Kolli N, Yergey A, et al. Nucleolar protein B23/nucleophosmin regulates the vertebrate SUMO pathway through SENP3 and SENP5 proteases. *J Cell Biol*. 2008 Nov 17;183(4):589-95.
- Yun JP, Chew EC, Liew CT, Chan JY, Jin ML, Ding MX, et al. Nucleophosmin/B23 is a proliferate shuttle protein associated with nuclear matrix. *J Cell Biochem*. 2003 Dec 15;90(6):1140-8.
- Yung BY. C-myc-mediated expression of nucleophosmin/B23 decreases during retinoic acid-induced differentiation of human leukemia HL-60 cells. *FEBS Lett*. 2004 Dec 17;578(3):211-6.
- Yung BY, Busch H, Chan PK. Translocation of nucleolar phosphoprotein B23 (37 kDa/pI 5.1) induced by selective inhibitors of ribosome synthesis. *Biochim Biophys Acta*. 1985 Dec 18;826(4):167-73.
- Zatsepina OV, Rousselet A, Chan PK, Olson MO, Jordan EG, Bornens M. The nucleolar phosphoprotein B23 redistributes in part to the spindle poles during mitosis. *J Cell Sci*. 1999 Feb;112 ( Pt 4)(Pt 4):455-66.
- Zeller KI, Haggerty TJ, Barrett JF, Guo Q, Wonsey DR, Dang CV. Characterization of nucleophosmin (B23) as a myc target by scanning chromatin immunoprecipitation. *J Biol Chem*. 2001 Dec 21;276(51):48285-91.
- Zhang W, Navenot JM, Frilot NM, Fujii N, Peiper SC. Association of nucleophosmin negatively regulates CXCR4-mediated G protein activation and chemotaxis. *Mol Pharmacol*. 2007 Nov;72(5):1310-21.
- Zhou BS. Tumour-initiating cells: Challenges and opportunities for anticancer drug discovery. *Nature reviews Drug discovery*. 2009;8(10).
- Zhu Y, Singh B, Hewitt S, Liu A, Gomez B, Wang A, et al. Expression patterns among interferon regulatory factor-1, human X-box binding protein-1, nuclear factor kappa B, nucleophosmin, estrogen receptor-alpha and progesterone receptor proteins in breast cancer tissue microarrays. *Int J Oncol*. 2006 Jan;28(1):67-76.

Zou Y, Wu J, Giannone RJ, Boucher L, Du H, Huang Y, et al. Nucleophosmin/B23 negatively regulates GCN5-dependent histone acetylation and transactivation. *J Biol Chem*. 2008 Feb 29;283(9):5728-37.

Zyss D, Gergely F. Centrosome function in cancer: Guilty or innocent? *Trends Cell Biol*. 2009 Jul;19(7):334-46.